

ON OPTIMAL RECOVERY OF PERIODIC ANALYTIC FUNCTIONS

K. YU. OSIPENKO, MOSCOW, RUSSIA

Let $S_\beta := \{z \in \mathbb{C} : |\operatorname{Im} z| < \beta\}$ be a strip in the complex plane. Denote by H_p^β the class of 2π -periodic, analytic in S_β functions f , which satisfy

$$\begin{aligned} \sup_{0 \leq \eta < \beta} \frac{1}{4\pi} \int_{\mathbb{T}} (|f(t + i\eta)|^p + |f(t - i\eta)|^p) dt &\leq 1, \quad 1 \leq p < \infty, \\ \sup_{z \in S_\beta} |f(z)| &\leq 1, \quad p = \infty. \end{aligned}$$

We consider the problem of optimal recovery of $Lf = f(\xi)$ or $f'(\xi)$, $\xi \in \mathbb{T}$, using the information $If = (f(x_1), \dots, f(x_n))$, $x_j \in \mathbb{T}$. We calculate the intrinsic error

$$e(L, H_p^\beta, I) := \inf_{A: \mathbb{C}^n \rightarrow \mathbb{C}} \sup_{f \in H_p^\beta} |Lf - A(If)| \tag{1}$$

and find an optimal algorithm A^* for which the infimum in (1) is attained.

For example, if $Lf = f'(0)$ and $If = (f(-h), f(h))$, then an optimal algorithm is given by

$$f'(0) \approx \frac{K}{\pi} \operatorname{dn} \frac{2(p-1)}{p} \frac{K}{\pi} h \frac{f(h) - f(-h)}{\operatorname{sn} \frac{2K}{\pi} h},$$

where $\operatorname{dn} z$ and $\operatorname{sn} z$ are the Jacobi elliptic functions with modulus k defined by the equation

$$\frac{\pi K'}{2K} = \beta$$

(K and K' are the complete elliptic integrals of the first kind with moduli k and $k' = \sqrt{1 - k^2}$).