

ISMAGILOV TYPE THEOREMS FOR LINEAR, GEL'FAND AND BERNSTEIN n -WIDTHS

K. YU. OSIPENKO AND O. G. PARFENOV

ABSTRACT. Using a variational principle for s -numbers, we obtain estimates for the linear, Gel'fand and Bernstein n -widths. A simple proof of some results concerned with the exact values of n -widths of diagonal operators is given. We also calculate the exact values of the Bernstein n -widths for the Hardy–Sobolev classes.

1. INTRODUCTION

Let X, Y be normed linear spaces and $T: X \rightarrow Y$ be a bounded linear operator. The linear λ_n , Gel'fand d^n and Bernstein b_n n -widths of the operator T are defined by

$$\begin{aligned}\lambda_n(T) &:= \inf_{P_n} \sup_{x \in BX} \|Tx - P_n x\|_Y, & d^n(T) &:= \inf_{X^n} \sup_{x \in BX^n} \|Tx\|_Y, \\ b_n(T) &:= \sup_{X_{n+1}} \inf_{\substack{x \in X_{n+1} \\ x \neq 0}} \frac{\|Tx\|_Y}{\|x\|_X},\end{aligned}$$

where P_n is any linear operator mapping X into Y of rank at most n , BX is the closed unit ball of X , X^n runs over all n -codimensional subspaces of X and X_{n+1} runs over all $(n+1)$ -dimensional subspaces of X .

In Osipenko and Stessin [1] the exact values of the linear and Gel'fand n -widths of the Hardy classes H_2 were obtained. A method of the proof was very close to the one from Ismagilov's Theorem [2] (see also [3, p.93]). After the paper [1] several results were obtained for $\lambda_n(T)$ and $d^n(T)$ where T is a map from a Hilbert space H into $C(E)$ (see [4]–[6]). Parfenov [7] solved an analogous problem for the Bernstein n -widths $b_n(T)$ where $T: L_\infty(E, \nu) \rightarrow H$ and ν is a probability measure on E .

In this paper we show that many of these results can be obtained, using a general principle concerned with extremal properties of s -numbers. Section 2 is devoted to this principle. In Section 3 we prove the estimates of the linear, Gel'fand and Bernstein n -widths. In Section 4 we give a simple proof of two results about the exact values of n -widths for diagonal operators in the discrete case. Finally, in Section 5 we calculate the Bernstein n -widths of the Hardy–Sobolev classes.

The research of the first author was supported in part by Russian Foundation of Fundamental Research and by the International Scientific Fund

The research of the second author was supported in part by the International Scientific Fund and by the Natural Sciences Academy of Russian Federation

2. VARIATIONAL PRINCIPLE FOR s -NUMBERS

Let H and H_1 be Hilbert spaces and $T: H \rightarrow H_1$ a bounded linear operator. Suppose that

$$T'T\varphi_k = \lambda_k\varphi_k, \quad k = 1, 2, \dots,$$

where $\lambda_1 \geq \lambda_2 \geq \dots > 0$ and $\{\varphi_k\}$ form a complete orthonormal basis for the range of $T'T$ (a sufficient condition is that T be a compact operator). The values $s_k(T) = \sqrt{\lambda_k}$ are called the s -numbers of T .

Set $\psi_k := s_k^{-1}(T)T\varphi_k$. Note that $\{\psi_k\}$ is an orthonormal system in H_1 . Then there exists the Schmidt decomposition of T (see, for example, [8]) which is given by

$$T = \sum_{k=1}^{\infty} s_k(T)(\cdot, \varphi_k)\psi_k.$$

Theorem 1. *Let T be as above. Then*

$$(1) \quad \sum_{k=1}^n s_k^2(T) = \max_{\{e_k\}_1^n} \sum_{k=1}^n \|Te_k\|_{H_1}^2$$

where the maximum is taken over all orthonormal systems $\{e_k\}_1^n$ in H . Furthermore,

$$(2) \quad \sum_{k=n+1}^{\infty} s_k^2(T) = \min_{\{e_k\}_1^{\infty}} \sum_{k=1}^{\infty} \|Te_k\|_{H_1}^2$$

where the minimum is taken over all orthonormal systems $\{e_k\}_1^{\infty}$ in H such that $\text{codim span}\{e_k\}_1^{\infty} \leq n$.

For a compact operator T this theorem was proved in Parfenov [6]. In our case the proof is almost the same because it does not so much depend on the compactness of T as on the fact that the eigenvectors $\{\varphi_k\}$ of $T'T$ form an orthonormal basis for the range of $T'T$.

We remark that both parts in (2) are finite iff

$$\sum_{k=1}^{\infty} s_k^2(T) < \infty.$$

In Parfenov [6] Theorem 1 was the basic tool in calculating the Gel'fand n -widths of operators $T: H \rightarrow L_{\infty}(E, \nu)$. Similar results were obtained in Osipenko [5], using Ismagilov's Theorem and the duality between the Kolmogorov and Gel'fand n -widths.

In order to estimate the Bernstein n -widths we need the following properties of s -numbers.

Theorem 2. *Let T be as in Theorem 1 and $\text{Ker } T = 0$. Then*

$$(3) \quad \sum_{k=1}^n s_k^{-2}(T) = \min_{\{f_k\}_1^n} \sum_{k=1}^n \|T^{-1}f_k\|_H^2$$

where the minimum is taken over all orthonormal systems $\{f_k\}_1^n$ in $T(H)$. Furthermore, if $\dim H = N < \infty$, then

$$(4) \quad \sum_{k=n+1}^N s_k^{-2}(T) = \max_{\{f_k\}_1^{N-n}} \sum_{k=1}^{N-n} \|T^{-1}f_k\|_H^2$$

where the maximum is taken over all orthonormal systems $\{f_k\}_1^{N-n}$ in $T(H)$.

Proof. Let $\{f_k\}_1^n$ be any orthonormal system in $T(H)$. Set $L_n := \text{span}\{T^{-1}f_k\}_1^n$, $\tilde{L}_n := \text{span}\{f_k\}_1^n$, and $T_n := T|_{L_n}$. Suppose that the Schmidt decomposition of T_n has the form

$$T_n = \sum_{k=1}^n s_k(T_n)(\cdot, \varphi'_k)\psi'_k.$$

The value

$$\left(\sum_{k=1}^n \|T_n^{-1}f_k\|_H^2 \right)^{1/2}$$

is the Hilbert–Schmidt norm of T_n^{-1} and does not depend on the choice of the orthonormal basis in \tilde{L}_n . Therefore

$$\sum_{k=1}^n \|T^{-1}f_k\|_H^2 = \sum_{k=1}^n \|T_n^{-1}f_k\|_H^2 = \sum_{k=1}^n \|T_n^{-1}\psi'_k\|_H^2 = \sum_{k=1}^n s_k^{-2}(T_n).$$

Let P_n be an orthoprojector in H onto L_n . Using the properties of s -numbers of bounded linear operators (see [8, p.82]) we have

$$s_k(T_n) = s_k(T \circ P_n) \leq \|P_n\| s_k(T) = s_k(T).$$

Thus

$$\sum_{k=1}^n \|T^{-1}f_k\|_H^2 \geq \sum_{k=1}^n s_k^{-2}(T).$$

If $f_k = \psi_k$, $k = 1, \dots, n$, then

$$\sum_{k=1}^n \|T^{-1}\psi_k\|_H^2 = \sum_{k=1}^n s_k^{-2}(T).$$

The equality (3) is proved.

To prove (4) note that

$$s_k(T^{-1}) = s_{N-k+1}^{-1}(T), \quad k = 1, \dots, N.$$

Now (4) follows from (1). The theorem is proved.

3. ESTIMATES OF LINEAR, GEL'FAND AND BERNSTEIN n -WIDTHS

To obtain estimates of n -widths we need the following simple result.

Lemma 1. *Let H be a Hilbert space, $\omega := \dim H$ and $T: H \rightarrow C(E)$ a bounded linear operator. Then*

$$\|T\|_{H \rightarrow C(E)} = \sup_{z \in E} \left(\sum_{k=1}^{\omega} |(Te_k)(z)|^2 \right)^{1/2}$$

for any orthonormal basis $\{e_k\}_1^{\omega}$ in H .

Proof. We have

$$\begin{aligned} \|T\|_{H \rightarrow C(E)} &= \sup_{h \in BH} \sup_{z \in E} |(Th)(z)| = \sup_{z \in E} \sup_{h \in BH} |(Th)(z)| \\ &= \sup_{z \in E} \sup_{\{c_k\}_1^{\omega} \in Bl_2} \left| \sum_{k=1}^{\omega} c_k (Te_k)(z) \right| = \sup_{z \in E} \left(\sum_{k=1}^{\omega} |(Te_k)(z)|^2 \right)^{1/2}. \end{aligned}$$

The lemma is proved.

Let H be a Hilbert space of functions defined on some set Ω . A function $K(z, w)$ defined on $\Omega \times \Omega$ is called a reproducing kernel of H if for each $w \in \Omega$, $K(z, w) \in H$ and for all $f \in H$

$$f(w) = (f(\cdot), K(\cdot, w))_H.$$

It is a well-known fact that if the $\{\varphi_k\}_1^{\omega}$ form an orthonormal basis in H , then

$$K(z, w) = \sum_{k=1}^{\omega} \varphi_k(z) \overline{\varphi_k(w)}.$$

Suppose that Ω is a topological space, $E \subset \Omega$ and $Tf := f|_E$ is a bounded linear operator from H into $C(E)$. Then from Lemma 1 we obtain

$$\|T\|_{H \rightarrow C(E)} = \sup_{z \in E} (K(z, z))^{1/2}.$$

Theorem 3 ([5], [6]). *Let H be a Hilbert space, E a topological space with probability measure ν such that $\text{supp } \nu = E$, and $T: H \rightarrow C(E)$ a bounded linear operator. Define $T_0: H \rightarrow L_2(E, \nu)$ by the equation $T_0 h := Th$. Assume that*

$$(5) \quad T_0' T_0 \varphi_k = s_k^2 \varphi_k, \quad k = 1, 2, \dots,$$

where $s_1 \geq s_2 \geq \dots > 0$ and $\{\varphi_k\}$ is an orthonormal basis for the range of $T_0' T_0$. Then

$$\left(\sum_{k=n+1}^{\infty} s_k^2 \right)^{1/2} \leq \lambda_n(T) = d^n(T) \leq \sup_{z \in E} \left(\sum_{k=n+1}^{\infty} |(T\varphi_k)(z)|^2 \right)^{1/2}.$$

Proof. Since $\text{supp } \nu = E$, $\text{Ker } T'_0 T_0 = \text{Ker } T_0 = \text{Ker } T$ and we can assume, without loss of generality, that $\{\varphi_k\}$ is an orthonormal basis in H . From the definition of the Gel'fand n -width it follows that

$$d^n(T) = \inf_{H^n} \|T\|_{H^n \rightarrow C(E)}$$

where H^n runs over all n -codimensional subspaces of H . Consider $H^n = \{\varphi_k\}_{n+1}^\infty$. We obtain from Lemma 1

$$d^n(T) \leq \sup_{z \in E} \left(\sum_{k=n+1}^{\infty} |(T\varphi_k)(z)|^2 \right)^{1/2}.$$

Let H^n be any n -codimensional subspace of H . Suppose that $\{\varphi'_k\}$ is an orthonormal basis in H^n . Using Lemma 1 and (2), we have

$$\begin{aligned} \|T\|_{H^n \rightarrow C(E)}^2 &= \sup_{z \in E} \sum_{k=1}^{\infty} |(T\varphi'_k)(z)|^2 \geq \int_E \sum_{k=1}^{\infty} |(T\varphi'_k)(z)|^2 d\nu(z) \\ &= \sum_{k=1}^{\infty} \|T_0 \varphi'_k\|_{L_2(E, \nu)}^2 \geq \sum_{k=n+1}^{\infty} s_k^2. \end{aligned}$$

Thus

$$d^n(T) \geq \left(\sum_{k=n+1}^{\infty} s_k^2 \right)^{1/2}.$$

The equality $\lambda_n(T) = d^n(T)$ is the well-known fact for operators defined on Hilbert spaces (see, [3, p.33]). The theorem is proved.

Now we will obtain the similar estimates for the Bernstein n -widths.

Theorem 4. *Let H , E and ν be as above, H_1 be a subspace of $L_2(E, \nu)$ and $X_E \subset H_1$ be a subspace of $C(E)$. Assume that a bounded linear operator $T_0: H_1 \rightarrow H$ satisfies the conditions (5) where $s_1 \geq s_2 \geq \dots > 0$, $\{\varphi_k\}$ is an orthonormal basis for the range of $T'_0 T_0$ and $\varphi_k \in X_E$, $k = 1, 2, \dots$. Define $T: X_E \rightarrow H$ by the equation $Tf := T_0 f$. Then*

$$(6) \quad \left(\sup_{z \in E} \sum_{k=1}^{n+1} s_k^{-2} |\varphi_k(z)|^2 \right)^{-1/2} \leq b_n(T) \leq \left(\sum_{k=1}^{n+1} s_k^{-2} \right)^{-1/2}.$$

Proof. Let $L_{n+1} \subset X_E$ and $\dim L_{n+1} = n+1$. Consider the operator $T_{n+1} := T|_{L_{n+1}}$. If $\text{Ker } T_{n+1} \neq 0$, then

$$\inf_{\substack{f \in L_{n+1} \\ f \neq 0}} \frac{\|Tf\|_H}{\|f\|_{C(E)}} = 0.$$

Suppose that $\text{Ker } T_{n+1} = 0$. Then we can define the operator $T_{n+1}^{-1}: T(L_{n+1}) \rightarrow L_{n+1}$. Using Lemma 1 and (4), for any orthonormal system $\{e_k\}_1^{n+1}$ in $T(L_{n+1})$ we have

$$\begin{aligned} \inf_{\substack{f \in L_{n+1} \\ f \neq 0}} \frac{\|Tf\|_H}{\|f\|_{C(E)}} &= \inf_{\substack{g \in T(L_{n+1}) \\ g \neq 0}} \frac{\|g\|_H}{\|T_{n+1}^{-1}g\|_{C(E)}} = \|T_{n+1}^{-1}\|_{T(L_{n+1}) \rightarrow C(E)}^{-1} \\ &= \left(\sup_{z \in E} \sum_{k=1}^{n+1} |(T_{n+1}^{-1}e_k)(z)|^2 \right)^{-1/2} \leq \left(\sum_{k=1}^{n+1} \int_E |(T_{n+1}^{-1}e_k)(z)|^2 d\nu(z) \right)^{-1/2} \\ &= \left(\sum_{k=1}^{n+1} \|T_{n+1}^{-1}e_k\|_{H_1}^2 \right)^{-1/2} \leq \left(\sum_{k=1}^{n+1} s_k^{-2}(T_{n+1}) \right)^{-1/2}. \end{aligned}$$

Since $s_k(T_{n+1}) \leq s_k(T_0) = s_k$, $k = 1, \dots, n+1$, we obtain

$$b_n(T) \leq \left(\sum_{k=1}^{n+1} s_k^{-2} \right)^{-1/2}.$$

Let $L_{n+1} = \text{span}\{\varphi_k\}_1^{n+1}$. Then $\psi_k := s_k^{-1}T\varphi_k$, $k = 1, \dots, n+1$, form an orthonormal system in $T(L_{n+1})$. Thus

$$\begin{aligned} b_n(T) &\geq \inf_{\substack{f \in L_{n+1} \\ f \neq 0}} \frac{\|Tf\|_H}{\|f\|_{C(E)}} = \left(\sup_{z \in E} \sum_{k=1}^{n+1} |(T_{n+1}^{-1}\psi_k)(z)|^2 \right)^{-1/2} \\ &= \left(\sup_{z \in E} \sum_{k=1}^{n+1} s_k^{-2} |\varphi_k(z)|^2 \right)^{-1/2}. \end{aligned}$$

The theorem is proved.

4. n -WIDTHS OF DIAGONAL OPERATORS

Let $T: l_2 \rightarrow l_\infty$ be the diagonal operator

$$(7) \quad T(\{x_k\}_1^\infty) := \{\lambda_k x_k\}_1^\infty$$

where $\lambda_1 \geq \lambda_2 \geq \dots > 0$. Smolyak [9] (in the finite-dimensional case) proved that

$$(8) \quad d^n(T) = \sup_{m > n} \left(\frac{m-n}{\sum_{k=1}^m \lambda_k^{-2}} \right)^{1/2}.$$

In dual terms this result was obtained by Sofman [10] (see also [11]). We will show that the lower bound in (8) easily follows from Theorem 3.

Denote by $\{e_k\}_1^\infty$ the standard basis of l_2 . Fix any $m > n$. Let $T_m: l_2^m \rightarrow l_\infty^m$ be the operator defined by

$$T_m(\{x_k\}_1^m) := \{\lambda_k x_k\}_1^m.$$

It is easy to see that $d^n(T) \geq d^n(T_m)$. Define the probability measure ν_m on the set $\{1, 2, \dots, m\}$ as follows

$$\nu_m(\{j\}) := \lambda_j^{-2} \left(\sum_{k=1}^m \lambda_k^{-2} \right)^{-1}.$$

Denote by T_{m0} the operator T_m regarded as an operator from l_2^m into $l_2^m(\nu_m)$. Then

$$T'_{m0} T_{m0} e_j = \left(\sum_{k=1}^m \lambda_k^{-2} \right)^{-1} e_j, \quad j = 1, \dots, m.$$

From Theorem 3

$$d^n(T_m) \geq \left(\frac{m-n}{\sum_{k=1}^m \lambda_k^{-2}} \right)^{1/2}.$$

Thus

$$\lambda_n(T) = d^n(T) \geq \sup_{m>n} d^n(T_m) \geq \left(\frac{m-n}{\sum_{k=1}^m \lambda_k^{-2}} \right)^{1/2}.$$

Note that the values (8) are also related to linear stochastic n -widths (see [12]).

Consider the operator (7) as an operator from l_∞ into l_2 . Here we assume that $\{\lambda_k\}_1^\infty \in l_2$. Galeev [13] proved the equality

$$b_n(T) = \min_{0 \leq m < n+1} \left(\frac{\sum_{k=m+1}^\infty \lambda_k^2}{n-m+1} \right)^{1/2}.$$

We will show how the upper bound can be obtained from Theorem 4.

Let $0 < \varepsilon < 1$ and $0 \leq m < n+1$. Define the probability measure ν_m on \mathbb{N} as follows

$$\nu_m(\{j\}) := \begin{cases} (1-\varepsilon) \frac{\lambda_j^2}{\sum_{k=m+1}^\infty \lambda_k^2}, & j > m \\ \frac{\varepsilon}{m}, & j \leq m. \end{cases}$$

Denote $T_0: l_2(\nu_m) \rightarrow l_2$ by the equation $T_0 x := T x$. It is easy to obtain that

$$T'_0 T_0 e_j = s_j^2 e_j, \quad j = 1, 2, \dots,$$

where

$$s_j^2 = \begin{cases} (1-\varepsilon)^{-1} \sum_{k=m+1}^\infty \lambda_k^2, & j > m \\ \varepsilon^{-1} \lambda_j^2 m, & j \leq m. \end{cases}$$

From Theorem 4

$$b_n(T) \leq \left((1-\varepsilon) \frac{n-m+1}{\sum_{k=m+1}^\infty \lambda_k^2} + \frac{\varepsilon}{m} \sum_{j=1}^m \lambda_j^{-2} \right)^{-1/2}.$$

Letting $\varepsilon \rightarrow 0$, we obtain

$$b_n(T) \leq \min_{0 \leq m < n+1} \left(\frac{\sum_{k=m+1}^\infty \lambda_k^2}{n-m+1} \right)^{1/2}.$$

5. BERNSTEIN n -WIDTHS OF HARDY-SOBOLEV CLASSES

Let A be a closed, convex centrally symmetric subset of a normed linear space Y . The Bernstein n -width of A is defined by

$$b_n(A, Y) := \sup_{Y_{n+1}} \sup \{ \lambda : \lambda B Y_{n+1} \subset A \}$$

where Y_{n+1} is any $(n+1)$ -dimensional subspace of Y . If $\text{Ker } T = 0$, then it is easily shown that

$$b_n(T) = b_n(T(BX), Y).$$

We need the following simple property of b_n .

Lemma 2. *Let H be a Hilbert space, A a closed, convex centrally symmetric subset of H and H_r an r -dimensional subspace of H such that $A \perp H_r$. Then*

$$(9) \quad b_{n+r}(A + H_r, H) = b_n(A, H).$$

Proof. Assume that $H_{n+1} \subset H$, $\dim H_{n+1} = n+1$ and

$$\sup \{ \lambda : \lambda B H_{n+1} \subset A \} = \mu > 0.$$

Put $H_{n+r+1} := H_{n+1} + H_r$. Since $A \perp H_r$ it follows that $H_{n+1} \perp H_r$ and $\dim H_{n+r+1} = n+r+1$. If $x \in H_{n+r+1}$, $\|x\|_H \leq \mu$, then $x = x_1 + x_2$ where $x_1 \in H_{n+1}$, $x_2 \in H_r$ and

$$\|x_1\|_H^2 \leq \|x_1\|_H^2 + \|x_2\|_H^2 = \|x\|_H^2 \leq \mu^2.$$

Thus $x_1 \in A$. Consequently $x \in A + H_r$. We have

$$\sup \{ \lambda : \lambda B H_{n+r+1} \subset A + H_r \} \geq \mu.$$

So we proved that

$$(10) \quad b_{n+r}(A + H_r, H) \geq b_n(A, H).$$

If $b_{n+r}(A + H_r, H) = 0$, then (9) follows from (10). Suppose that $b_{n+r}(A + H_r, H) > 0$. Let $H_{n+r+1} \subset H$, $\dim H_{n+r+1} = n+r+1$ and

$$\sup \{ \lambda : \lambda B H_{n+r+1} \subset A + H_r \} = \mu > 0.$$

Since $H_{n+r+1} \subset \text{span } A + H_r$, $\dim(H_{n+r+1} \cap \text{span } A) \geq n+1$. Hence there exists a subspace $H_{n+1} \subset H_{n+r+1} \cap \text{span } A$ with $\dim H_{n+1} = n+1$. Let $x \in H_{n+1}$ and $\|x\|_H \leq \mu$. Then $x \in A + H_r$. In addition $x \in \text{span } A$. Thus $x \in A$ and

$$\sup \{ \lambda : \lambda B H_{n+1} \subset A \} \geq \mu.$$

Therefore

$$b_n(A, H) \geq b_{n+r}(A + H_r, H).$$

The lemma is proved.

Theorem 5. Let H , E , ν , H_1 and X_E be as in Theorem 4. Suppose that the Hilbert space H_1 has a reproducing kernel. Let $\{\varphi_k\}$ be an orthonormal basis in H_1 and $T_0: H_1 \rightarrow H$ a bounded linear operator. Assume that $\{T_0\varphi_k\}$ is an orthogonal system in H , $\varphi_k \in X_E$, $k = 1, 2, \dots$ and $s_k := \|T_0\varphi_k\|_H$ form a non-increasing sequence. Define $T: X_E \rightarrow H$ by the equation $Tf := T_0f$. Then the inequalities (6) hold.

Proof. Denote by $K(z, w)$ the reproducing kernel of H_1 . Since $\{\varphi_k\}$ is an orthonormal basis in H_1 the representation

$$K(z, w) = \sum_{k=1}^{\infty} \varphi_k(z) \overline{\varphi_k(w)}$$

holds. We have

$$(T'_0 T_0 f)(w) = ((T'_0 T_0 f)(\cdot), K(\cdot, w))_{H_1} = ((T_0 f)(\cdot), (T_0 K)(\cdot, w))_H.$$

Thus

$$(T'_0 T_0 \varphi_j)(w) = ((T_0 \varphi_j)(\cdot), \sum_{k=1}^{\infty} \overline{\varphi_k(w)} (T_0 \varphi_k)(\cdot))_H = s_j^2 \varphi_j(w).$$

Now it suffices to apply Theorem 4. The theorem is proved.

Let B_ρ be the ball of \mathbb{C}^n of radius ρ

$$B_\rho := \{z := (z_1, \dots, z_n) \in \mathbb{C}^n : |z|^2 := \sum_{k=1}^n |z_k|^2 < \rho\},$$

$S_\rho := \partial B_\rho$, σ_ρ the probability measure on the sphere S_ρ which is invariant with respect to the orthogonal group $O(2n)$ and ν_ρ the normalized Lebesgue measure in B_ρ (if $\rho = 1$ we will write B , S , σ and ν).

The Hardy space $H_p(B)$ (H_p) is the set of holomorphic functions in B which satisfy

$$\|f\|_{H_p} := \sup_{0 < \rho < 1} \left(\int_S |f(z)|^p d\sigma(z) \right)^{1/p} < \infty, \quad 1 \leq p < \infty,$$

$$\|f\|_{H_\infty} := \sup_{z \in B} |f(z)|.$$

Let $f(z)$ be a holomorphic function in B and

$$f(z) = \sum_{k=0}^{\infty} F_k(z)$$

be a homogeneous decomposition of f . The radial derivative of order r is defined by

$$\mathcal{R}^r f(z) := \sum_{k=r}^{\infty} \frac{k!}{(k-r)!} F_k(z)$$

(for $r = 1$ see [14, Chap.6]). Denote by $H\mathcal{R}_\infty^r(B)$ ($H\mathcal{R}_\infty^r$) the class of holomorphic functions in B for which $\mathcal{R}^r f \in BH_\infty$.

Set

$$N_m := \sum_{k=0}^{m-1} \binom{n+k-1}{n-1}.$$

Note that $N_m = \dim \mathcal{P}_{m-1}^n$ where \mathcal{P}_m^n is the space of n -variable polynomials of degree m or less.

Theorem 6. *For all $0 < \rho \leq 1$ and all $m \geq r + 1$*

(11)

$$b_{N_m-1}(H\mathcal{R}_\infty^r, L_2(S_\rho, \sigma_\rho)) = \left(\frac{1}{(n-1)!} \sum_{k=r}^{m-1} \frac{k!(n+k-1)!}{((k-r)!)^2} \rho^{-2k} \right)^{-1/2},$$

(12)

$$b_{N_m-1}(H\mathcal{R}_\infty^r, L_2(B_\rho, \nu_\rho)) = \left(\frac{1}{n!} \sum_{k=r}^{m-1} \frac{k!(n+k)!}{((k-r)!)^2} \rho^{-2k} \right)^{-1/2}.$$

Proof. For multiindex $\alpha := (\alpha_1, \dots, \alpha_n)$ and $z \in \mathbb{C}^n$ set

$$z^\alpha := z_1^{\alpha_1} \dots z_n^{\alpha_n}, \quad |\alpha| := \alpha_1 + \dots + \alpha_n, \quad \alpha! := \alpha_1! \dots \alpha_n!,$$

$$D_j := \frac{\partial}{\partial z_j}, \quad D^\alpha := D_1^{\alpha_1} \dots D_n^{\alpha_n}.$$

Denote by H_p^0 the space of all functions $f \in H_p$ for which $(D^\alpha f)(0) = 0$, $|\alpha| = 0, \dots, r-1$. It is known (see [14]) that functions from H_p , $1 \leq p \leq \infty$, have finite boundary values almost everywhere. Moreover, H_2 is a Hilbert space with the inner product

$$(f, g)_{H_2} := \int_S f(z) \overline{g(z)} d\sigma(z).$$

The space H_2 has the reproducing kernel

$$K(z, w) = \left(1 - \sum_{k=1}^n z_k \overline{w_k} \right)^{-n}.$$

Define $T_0: H_2^0 \rightarrow L_2(S_\rho, \sigma_\rho)$ and $T: H_\infty^0 \rightarrow L_2(S_\rho, \sigma_\rho)$ by the equations

$$(13) \quad (T_0 f)(z) := \sum_{k=r}^{\infty} \frac{(k-r)!}{k!} F_k(z), \quad T f := T_0 f,$$

where

$$f(z) = \sum_{k=r}^{\infty} F_k(z).$$

It is easy to see that

$$H\mathcal{R}_\infty^r = T(BH_\infty^0) + \mathcal{P}_{r-1}^n.$$

Since monomials z^α are orthogonal in $L_2(S_\rho, \sigma_\rho)$ we obtain from Lemma 2

$$(14) \quad b_{N_m-1}(H\mathcal{R}_\infty^r, L_2(S_\rho, \sigma_\rho)) = b_{N_m-N_r-1}(T).$$

For every $0 < \rho \leq 1$ (see [14])

$$\|z^\alpha\|_{L_2(S_\rho, \sigma_\rho)}^2 = \frac{(n-1)!\alpha!}{(n+|\alpha|-1)!} \rho^{2|\alpha|}.$$

Thus the functions

$$\varphi_\alpha(z) := \left(\frac{(n+|\alpha|-1)!}{(n-1)!\alpha!} \right)^{1/2} z^\alpha, \quad |\alpha| \geq r,$$

form a complete orthonormal basis in H_2^0 . We have

$$\|T_0\varphi_\alpha\|_{L_2(S_\rho, \sigma_\rho)}^2 = \left(\frac{(|\alpha|-r)!}{|\alpha|!} \right)^2 \rho^{2|\alpha|}.$$

The number of different monomials z^α with $|\alpha| = k$ equals $\binom{n+k-1}{n-1}$. By

Theorem 5

$$\begin{aligned} \left(\sup_{z \in S} \sum_{|\alpha|=r}^{m-1} \left(\frac{|\alpha|!}{(|\alpha|-r)!} \right)^2 \rho^{-2|\alpha|} |\varphi_\alpha(z)|^2 \right)^{-1/2} &\leq b_{N_m-N_r-1}(T) \\ &\leq \left(\sum_{k=r}^{m-1} \left(\frac{k!}{(k-r)!} \right)^2 \binom{n+k-1}{n-1} \rho^{-2k} \right)^{-1/2}. \end{aligned}$$

Using the equation

$$\sum_{|\alpha|=k} \frac{|z^{2\alpha}|}{\alpha!} = \frac{|z|^{2k}}{k!},$$

we obtain

$$b_{N_m-N_r-1}(T) = \left(\frac{1}{(n-1)!} \sum_{k=r}^{m-1} \frac{k!(n+k-1)!}{((k-r)!)^2} \rho^{-2k} \right)^{-1/2}.$$

Now (11) follows from (14).

The proof of (12) is almost the same. The difference is that we have to consider the operators $T_0: H_2^0 \rightarrow L_2(B_\rho, \nu_\rho)$ and $T: H_\infty^0 \rightarrow L_2(B_\rho, \nu_\rho)$ defined by (13). Then we use

$$\|T_0\varphi_\alpha\|_{L_2(B_\rho, \nu_\rho)}^2 = \left(\frac{(|\alpha|-r)!}{|\alpha|!} \right)^2 \frac{n}{n+|\alpha|} \rho^{2|\alpha|}.$$

The theorem is proved.

For $n = 1$ the class $H\mathcal{R}_\infty^r$ coincides with the class BH_∞^r defined as the set of all holomorphic functions in B for which $f^{(r)}(z) \in BH_\infty$. From Theorem 6 we obtain the following result.

Corollary 1. *For all $0 < \rho \leq 1$ and all $m \geq r$*

$$b_m(BH_\infty^r, L_2(S_\rho, \sigma_\rho)) = \left(\sum_{k=r}^m \left(\frac{k!}{(k-r)!} \right)^2 \rho^{-2k} \right)^{-1/2},$$

$$b_m(BH_\infty^r, L_2(B_\rho, \nu_\rho)) = \left(\sum_{k=r}^m \frac{k!(k+1)!}{((k-r)!)^2} \rho^{-2k} \right)^{-1/2}.$$

In particular, we have for $r = 0$, $m \geq 0$ and $0 < \rho < 1$

$$(15) \quad b_m(BH_\infty, L_2(S_\rho, \sigma_\rho)) = \rho^m \left(\frac{1 - \rho^2}{1 - \rho^{2m+2}} \right)^{1/2},$$

$$(16) \quad b_m(BH_\infty, L_2(B_\rho, \nu_\rho)) = \rho^m \frac{1 - \rho^2}{\sqrt{(m+1)(1 - \rho^2) - \rho^2(1 - \rho^{2m+2})}}.$$

The values (11) for $r = 0$ and (15) were calculated in [7].

Let us compare (15) and (16) with the exact values of the Kolmogorov, linear, and Gel'fand n -widths. From [15] and [16] it follows that

$$d_m(BH_\infty, X) = \lambda_m(BH_\infty, X) = d^m(BH_\infty, X) = \begin{cases} \rho^m, & X = L_2(S_\rho, \sigma_\rho) \\ \frac{\rho^m}{\sqrt{m+1}}, & X = L_2(B_\rho, \nu_\rho). \end{cases}$$

Finally, we will determine exact values of the Bernstein n -widths for some classes of periodic holomorphic functions. Let $D_\beta := \{z \in \mathbb{C} : |\operatorname{Im} z| < \beta\}$. Denote by $\tilde{H}_{p,\beta}$ the set of all 2π -periodic holomorphic functions in D_β which satisfy the conditions

$$\|f\|_{\tilde{H}_{p,\beta}} := \sup_{0 < h < \beta} \left(\frac{1}{4\pi} \int_0^{2\pi} (|f(x+ih)|^p + |f(x-ih)|^p) dx \right)^{1/p} < \infty, \quad 1 \leq p < \infty,$$

$$\|f\|_{\tilde{H}_{\infty,\beta}} := \sup_{z \in D_\beta} |f(z)| < \infty.$$

Let $B\tilde{H}_{p,\beta}^r$ be the set of all 2π -periodic holomorphic functions in D_β for which $f^{(r)}(z) \in B\tilde{H}_{p,\beta}$. Denote by L_2 the periodic complex-valued Lebesgue space on the real axis with the norm

$$\|f\|_{L_2} := \left(\frac{1}{2\pi} \int_0^{2\pi} |f(x)|^2 dx \right)^{1/2}.$$

Theorem 7.

(1) *For all $n \geq 1$ and $r \geq 1$*

$$b_{2n}(B\tilde{H}_{\infty,\beta}^r, L_2) = \left(2 \sum_{k=1}^n k^{2r} \cosh 2k\beta \right)^{-1/2}.$$

(2) *For all $n \geq 0$*

$$(17) \quad b_{2n}(B\tilde{H}_{\infty,\beta}, L_2) = \left(\frac{\sinh \beta}{\sinh(2n+1)\beta} \right)^{1/2}.$$

Proof. The space $\tilde{H}_{2,\beta}$ is the Hilbert space with the inner product

$$(f, g)_{\tilde{H}_{2,\beta}} := \frac{1}{4\pi} \int_0^{2\pi} \left(f(x + i\beta) \overline{g(x + i\beta)} + f(x - i\beta) \overline{g(x - i\beta)} \right) dx.$$

The functions

$$\varphi_k(z) := \frac{e^{ikz}}{\sqrt{\cosh 2k\beta}}, \quad k \in \mathbb{Z},$$

form an orthonormal basis in $\tilde{H}_{2,\beta}$. The space $\tilde{H}_{2,\beta}$ has the reproducing kernel

$$K(z, w) = \sum_{k \in \mathbb{Z}} \varphi_k(z) \overline{\varphi_k(w)} = 1 + 2 \sum_{k=1}^{\infty} \frac{\cos k(z - \overline{w})}{\cosh 2k\beta}.$$

Let $r \geq 1$. Denote by $\tilde{H}_{p,\beta}^0$ the space of functions $f \in \tilde{H}_{p,\beta}$ for which

$$\int_0^{2\pi} f(x) dx = 0.$$

Define $T_0: \tilde{H}_{2,\beta}^0 \rightarrow L_2$ and $T: \tilde{H}_{\infty,\beta}^0 \rightarrow L_2$ by the equations

$$(T_0 f)(z) := \sum_{\substack{k \in \mathbb{Z} \\ k \neq 0}} \frac{c_k}{(ik)^r} e^{ikz}, \quad T f := T_0 f,$$

where

$$f(z) = \sum_{\substack{k \in \mathbb{Z} \\ k \neq 0}} c_k e^{ikz}.$$

It is easily seen that

$$B\tilde{H}_{\infty,\beta}^r = T(B\tilde{H}_{\infty,\beta}^0) + \mathbb{C}.$$

By Lemma 2 we obtain

$$b_{2n}(B\tilde{H}_{\infty,\beta}^r, L_2) = b_{2n-1}(T).$$

The functions $\varphi_k(z)$, $k = \pm 1, \pm 2, \dots$ form a complete orthonormal basis in $\tilde{H}_{2,\beta}^0$ and

$$\|T_0 \varphi_k\|_{L_2}^2 = \frac{1}{k^{2r} \cosh 2k\beta}.$$

Since for all $z \in \partial D_\beta$

$$|\varphi_k(z)|^2 + |\varphi_{-k}(z)|^2 = 2$$

we have by Theorem 5

$$b_{2n-1}(T) = \left(2 \sum_{k=1}^n k^{2r} \cosh 2k\beta \right)^{-1/2}.$$

To obtain (2) we use the same scheme and the equality

$$1 + 2 \sum_{k=1}^n \cosh 2k\beta = \frac{\sinh(2n+1)\beta}{\sinh \beta}.$$

The theorem is proved.

Denote by $B\tilde{H}_{\infty,\beta}^{\mathbb{R}}$ the set of functions from $B\tilde{H}_{\infty,\beta}$ that are real-valued on \mathbb{R} . For even n the exact values of the Kolmogorov, linear, and Gel'fand n -widths of $B\tilde{H}_{\infty,\beta}^{\mathbb{R}}$ in L_q , $1 \leq q \leq \infty$, were determined in [17]. In particular, for $q = 2$

$$\begin{aligned} d_{2n}(B\tilde{H}_{\infty,\beta}^{\mathbb{R}}, L_2) &= \lambda_{2n}(B\tilde{H}_{\infty,\beta}^{\mathbb{R}}, L_2) = d^{2n}(B\tilde{H}_{\infty,\beta}^{\mathbb{R}}, L_2) \\ &= \left(\frac{\lambda}{\Lambda} \int_0^1 \frac{t^2 dt}{\sqrt{(1-t^2)(1-\lambda^2 t^2)}} \right)^{1/2} = \sqrt{2}e^{-\beta n} + O(e^{-5\beta n}), \end{aligned}$$

where Λ is the complete elliptic integral of the first kind with modulus

$$\lambda = 4e^{-2\beta n} \left(\sum_{k=0}^{\infty} e^{-4\beta n k(k+1)} \right)^2 \left(1 + 2 \sum_{k=1}^{\infty} e^{-4\beta n k^2} \right)^{-2}.$$

By Theorem 5 it can be shown that (17) also holds in the real case for the class $B\tilde{H}_{\infty,\beta}^{\mathbb{R}}$. Thus

$$b_{2n}(B\tilde{H}_{\infty,\beta}^{\mathbb{R}}, L_2) = \sqrt{1 - e^{-2\beta}} e^{-\beta n} + O(e^{-5\beta n}).$$

REFERENCES

1. K. Yu. Osipenko and M. I. Stessin, *On n -widths of the Hardy class H^2 in the unit ball of \mathbb{C}^n* , Uspekhi Mat. Nauk **45** (1990), 193–194; English transl. in Russian Math. Surveys **45** (1990), 235–236.
2. R. S. Ismagilov, *On n -dimensional diameters of compacts in a Hilbert space*, Funktsional. Anal. i Prilozhen. **2** (1968), 32–39; English transl. in Functional Anal. Appl. **2** (1968), 125–132.
3. A. Pinkus, *n -Widths in Approximation Theory*, Springer-Verlag, Berlin, 1980.
4. S. D. Fisher and M. I. Stessin, *On n -widths of classes of holomorphic functions with reproducing kernels*, Illinois J. Math. **38** (1994), 589–615.
5. K. Yu. Osipenko, *On N -widths of holomorphic functions of several variables*, J. Approx. Theory **82** (1995), 135–155.
6. O. G. Parfenov, *Gel'fand n -widths of some classes of analytic functions*, Mat. Sb. **185** (1994), 105–112. (Russian)
7. O. G. Parfenov, *Bernstein n -widths of some classes of analytic functions*, Mat. Zametki **56** (1994), 123–131. (Russian)
8. I. C. Gohberg and M. G. Krein, *Introduction to the Theory of Linear Nonselfadjoint Operators*, Nauka, Moscow, 1965; English transl. in Transl. Math. Monographs, vol. 18, Amer. Math. Soc., Providence, R. I., 1969.
9. S. A. Smolyak, *On optimal restoration of functions and functionals of them*, Candidate Dissertation, Moscow State University, Moscow, 1965. (Russian)
10. L. B. Sofman, *Diameters of octahedra*, Mat. Zametki **5** (1969), 429–436; English transl. in Math. Notes **5** (1969), 258–262.

11. C. V. Hutton, J. S. Morrell and J. R. Retherford, *Diagonal operators, approximation numbers, and Kolmogoroff diameters*, J. Approx. Theory **16** (1976), 48–80.
12. E. Novak, *Optimal linear randomized methods for linear operators in Hilbert spaces*, J. Complexity **8** (1992), 22–36.
13. E. M. Galeev, *Bernstein diameters of infinite dimensional ellipsoids*, in Complex Analysis. Differential and Integral Equations, Elista, 1990, pp. 14–20. (Russian)
14. W. Rudin, *Function Theory in the Unit Ball of \mathbb{C}^n* , Springer-Verlag, New-York, 1980.
15. O. G. Parfenov, *Gel'fand diameters of the unit ball of the Hardy class H^p in weight spaces*, Mat. Zametki **37** (1985), 171–175; English transl. in Math. Notes **37** (1985), 96–98.
16. Yu. A. Farkov, *Widths of Hardy and Bergman classes in a ball in \mathbb{C}^n* , Uspekhi Mat. Nauk **45** (1990), 197–198; English transl. in Russian Math. Surveys **45** (1990), 229–231.
17. K. Yu. Osipenko, *On the n -widths, optimal quadrature formulae and optimal recovery of functions which are analytic in a strip*, Izv. Ross. Akad. Nauk Ser. Mat. **58** (1994), 55–79. (Russian)

DEPARTMENT OF MATHEMATICS, MOSCOW STATE UNIVERSITY OF AVIATION TECHNOLOGY,
MOSCOW, RUSSIA, 103767

DEPARTMENT OF MATHEMATICS, PSKOV PEDAGOGICAL INSTITUTE, PSKOV, RUSSIA, 180000