

SCHWARZ LEMMA AND OPTIMAL RECOVERY OF FUNCTIONS IN H^2

Konstantin Yu. Osipenko,¹ Michael I. Stessin²

¹ "MATI" - Russian State Technological University, Orshanskaya Str. 3, Moscow, 121552, Russia

² Department of Mathematics and Statistics, University at Albany, Albany, NY 12222, USA

e-mail: ¹kosipenko@yahoo.com, ²stessin@math.albany.edu

AMS Subject Classification: 30H05, 30E10

Keywords. Extremal problems, optimal recovery, Hardy spaces, Shwartz lemma

Let $D \subset C^k$ be a domain, ν be a probability measure on \bar{D} and X be a closed subspace of $L^2(\nu)$. Consider $D_0, \dots, D_n \subset D$ and probability measures μ_0, \dots, μ_n on D_0, \dots, D_n respectively. We suppose that $X \subset L^2(\mu_j)$, $j = 0, 1, \dots, n$. We allow one of D_j to coincide with D . In this case we assume that μ_j coincides with ν .

Write $\mathcal{D} = (D_0, \dots, D_n)$, $\mu = (\mu_0, \dots, \mu_n)$, $\mu = (\mu_1, \dots, \mu_n)$, $y = (y_1, \dots, y_n)$.

1. OPTIMAL RECOVERY PROBLEM

Given y_1, \dots, y_n defined on D_1, \dots, D_n such that

$$\|f_j - y_j\|_{L^2(\mu_j)} \leq \delta_j, \quad j = 1, \dots, n,$$

we are to reconstruct f . Here f_j is the restriction of f to D_j and $\delta_j \geq 0$, $j = 1, \dots, n$ are accuracy levels. In particular, $\delta_j = 0$ means that f is known precisely on D_j .

A recovery algorithm (method, procedure, etc.) is an operator

$$A: L^2(\mu_1) \times \dots \times L^2(\mu_n) \rightarrow L^2(\mu_0).$$

We consider $A(y)$, $y = (y_1, \dots, y_n)$, to be the recovered value of f on D_0 . At this point we impose no conditions on A .

The maximal possible error of a method A is

$$e(X, \mathcal{D}, \mu, \delta, A) = \sup_{\substack{f \in X, y \in L^2(\mu_1) \times \dots \times L^2(\mu_n) \\ \|f_j - y_j\|_{L^2(\mu_j)} \leq \delta_j, j=1, \dots, n}} \|f_0 - A(y)\|_{L^2(\mu_0)}$$

The optimal recovery error is

$$E(X, \mathcal{D}, \mu, \delta) = \inf_{A: L^2(\mu_1) \times \dots \times L^2(\mu_n) \rightarrow L^2(\mu_0)} e(X, \mathcal{D}, \mu, \delta, A).$$

A method \hat{A} such that

$$E(X, \mathcal{D}, \mu, \delta) = e(X, \mathcal{D}, \mu, \delta, \hat{A})$$

is called an *optimal recovery method*.

The problem of finding an optimal recovery method (and sometimes an extremal function at which the optimal recovery error is attained) is usually referred to as *optimal recovery problem*.

2. EXTREMAL PROBLEM

The optimal recovery problem is closely related to the following extremal problem. Find

$$(1) \quad \|f_0\|_{L^2(\mu_0)} \rightarrow \max, \quad \|f_j\|_{L^2(\mu_j)}^2 \leq \delta_j^2, \quad j = 1, \dots, n, \quad f \in X.$$

A special case of this extremal problem is when D is the unit disk \mathbb{D} , μ_0 and μ_1 are point masses and μ_2 is the normalized Lebesgue measure on the unit circle. Here the problem turns into

$$\max\{|f(a_0)| : |f(a_1)| \leq \delta_1, \|f\|_{H^2} \leq \delta_2\},$$

which might be viewed as a version of the classical Schwarz lemma. Here we consider another variant of Schwarz Lemma. Let $a \in \mathbb{D}$ and Γ be a circle inside of the unit disk, μ be the normalized Lebesgue measure on Γ , and $\mu > 0$. Find

$$(2) \quad \sup \left\{ \int_{\Gamma} |f|^2 d\mu : f \in H^2, \|f\|_{H^2}^2 \leq 1, |f(a)| \leq \delta \right\}.$$

We will consider the case when the circle Γ passes through the origin and its center lies on the real axis, so that

$$\Gamma = \{z \in \mathbb{C} : |z - \rho| = \rho\}, \quad 0 < \rho < 1/2.$$

The corresponding optimal recovery problem is: *Reconstruct a Hardy function f on the circle Γ from its value at a given with some tolerance.*

There are several papers where similar problems were considered for Hardy and Bergman spaces in connection with optimal recovery in both one and several dimensional cases (see, for example, [4]–[6]).

3. EULER EQUATION FOR THE GENERAL PROBLEM

Let $K(z, w)$ be the reproducing kernel of X . Write

$$\tilde{\mu} = -\mu_0 + \sum_{j=1}^n \lambda_j \mu_j.$$

Then $\tilde{\mu}$ is a regular measure on D and every function from X is square-integrable with respect to $\tilde{\mu}$. For $w \in D$ we introduce

$$d\tilde{\mu}_w(z) = K(z, w)d\tilde{\mu}(z).$$

Obviously every function from X is $\tilde{\mu}_w$ -integrable.

We further define

$$\tau_w^\lambda(z) = \int_D K(z\tau)d\tilde{\mu}_w(\tau).$$

Theorem 1. *If $\tilde{f} \in X$ is a solution of the general extremal problem above, then there exists a non-negative vector $\hat{\lambda} = (\hat{\lambda}_1, \dots, \hat{\lambda}_n)$ such that*

$$\hat{f} = (\text{span}\{\tau_w^{\hat{\lambda}}, w \in D\})^\perp,$$

and

$$\hat{\lambda}_j(\|f\|_{L_2(\mu_j)} - \delta_j) = 0, j = 1, \dots, n.$$

We say that a non-negative vector $\lambda = (\lambda_1, \dots, \lambda_n)$ belongs to the spectrum of the problem, if there exists an admissible for this problem function $f \in X$ such that

1. $\lambda_j(\|f\|_{L_2(\mu_j)} - \mu_j) = 0$.
2. $f \in (\text{span}\{\tau_w^\lambda : w \in D\})^\perp$.

In this case we call f a *spectral function*.

Theorem 2. *Let Λ be the spectrum of the problem. Then*

$$(3) \quad \sup_{\substack{f \in X \\ \|f\|_{L_2(\mu_j)} \leq \delta_j, j=1, \dots, n}} = \sup_{\lambda \in \Lambda} \sum_{j=1}^n \lambda_j \delta_j^2.$$

We call a spectral point $(\hat{\lambda}_1, \dots, \hat{\lambda}_n)$ *extremal*, if the maximum of the right-hand side of (3) is attained at $(\hat{\lambda}_1, \dots, \hat{\lambda}_n)$.

4. SPECTRUM OF THE SCHWARZ LEMMA

Here we have.

$$\begin{aligned} \tau_w^\lambda &= -\frac{1}{\pi} \int_{\Gamma} \frac{1}{1-z\bar{\tau}} \cdot \frac{1}{1-\tau\bar{w}} \cdot \frac{|d\tau|}{|\tau-\rho|} + \\ &\lambda_1 \frac{1}{1-z\bar{a}} \cdot \frac{1}{1-a\bar{w}} + \frac{\lambda_2}{2\pi} \int_{|\tau|=1} \frac{1}{1-z\bar{\tau}} \cdot \frac{1}{1-\tau\bar{w}} |d\tau| = \\ &-\frac{1}{1-z\rho-\rho\bar{w}} + \frac{\lambda_1}{(1-z\bar{a})(1-a\bar{w})} + \frac{\lambda_2}{1-z\bar{w}}. \end{aligned}$$

By Theorem 1 every extremal function satisfies the following equation

$$\frac{1}{1-\rho w} f\left(\frac{\rho}{1-\rho w}\right) = \lambda_1 \frac{f(a)}{1-\bar{a}w}$$

for some $\lambda_1, \lambda_2 \geq 0$ and all $w \in \mathbb{D}$. Let

$$b = \frac{1 - \sqrt{1-4\rho^2}}{2\rho}.$$

Then b is the Denjoy-Wolff point of the following self-mapping of \mathbb{D}

$$z \rightarrow \frac{\rho}{1-\rho z},$$

and the disk bounded by the circle Γ is a hyperbolic neighborhood of b .

Consider the following functions

$$\varphi_j(z) = \frac{\sqrt{1-b^2}}{1-bz} \left(\frac{b-z}{1-bz}\right)^j, \quad j = 0, 1, \dots$$

These functions form an orthonormal basis of H^2 , and they are eigenfunctions of the operator

$$Tf(z) = \frac{1}{1-\rho z} f\left(\frac{\rho}{1-\rho z}\right),$$

and the corresponding eigenvalues are

$$\alpha_j = \frac{b^{2j}}{1-\rho b}.$$

Theorem 3. *Let $a \neq b$.*

1. *If*

$$\left|a - \frac{\rho}{1-\rho^2}\right| \geq \frac{\rho^2}{1-\rho^2},$$

or

$$\delta > \frac{\sqrt{|a|^2\rho^2 - |\rho - a|^2}}{a\rho + \bar{a}\rho - |a|^2},$$

then the spectrum of Schwarz Lemma extremal problem consists of two parts $\Lambda = \Lambda_1 \cup \Lambda_2$, where

$$\Lambda_1 = \{(0, \alpha_j) : |\varphi_j(a)| \leq \delta\},$$

$$\Lambda_2 = \{(\lambda_1, \lambda_2) : \lambda_1, \lambda_2 > 0, F(\lambda_2) = \delta^{-2}, \lambda_1 = h(\lambda_2)\},$$

where

$$F(\lambda) = \sum_{j=0}^{\infty} \frac{|\varphi_j(a)|^2}{(a_j - \lambda)^2} h^2(\lambda), \quad h(\lambda) = \left(\sum_{j=0}^{\infty} \frac{|\varphi_j(a)|^2}{a_j - \lambda} \right)^{-1}.$$

2. If

$$\left| a - \frac{\rho}{1 - \rho^2} \right| < \frac{\rho^2}{1 - \rho^2},$$

and

$$\delta \leq \frac{\sqrt{|a|^2 \rho^2 - |\rho - a|^2}}{a\rho + \bar{a}\rho - |a|^2},$$

then the spectrum includes in addition the point

$$\Lambda_3 = \left\{ \left(\frac{a\rho + \bar{a}\rho - |a|^2}{\rho^2}, 0 \right) \right\}.$$

Theorem 4. Let $a = b$,

$$\Lambda_1 = \{(0, \alpha_j) : j = 1, 2, \dots, \},$$

$$\Lambda_2 = \{((1 - b^2)(\alpha_0 - \alpha_j), \alpha_j) : j = 1, 2, \dots, \}.$$

Then the spectrum of problem is $\Lambda = \Lambda_1 \cup \Lambda_2$, if $\delta < \frac{1}{\sqrt{1 - b^2}}$, and

$$\Lambda = \Lambda_1 \cup \Lambda_2 \cup \{(0, \alpha_0)\}, \text{ if } \delta \geq \frac{1}{\sqrt{1 - b^2}}.$$

It turns out that Λ_2 is the most important part of the spectrum.

Proposition 1. If a lies outside Γ , then $F(\lambda) \rightarrow \infty$ as $\lambda \rightarrow 0$.

This Proposition implies that if a lies outside Γ , then Λ_2 contains only finite number of points.

Now we will use Theorem 2 to describe the extremal points of the spectrum.

Proposition 2. If $\delta \geq |\varphi_0(a)|$, then $(0, \alpha_0)$ is the extremal point of the spectrum.

Proposition 3. If $a = b$ and $\delta < 1/\sqrt{1 - b^2}$, then the extremal spectral point is

$$(\widehat{\lambda}_1, \widehat{\lambda}_2) = ((1 - b^2)(\alpha_0 - \alpha_1), \alpha_1).$$

Proposition 4. If $\delta < |\varphi_0(a)|$, then Λ_1 does not contain extremal spectral points.

Note that the function

$$g(\lambda) = \sum_{j=0}^{\infty} \frac{|\varphi_j(a)|^2}{\alpha_j - \lambda}$$

is monotone and increases from $-\infty$ to $+\infty$ when $\lambda \in (\alpha_{j+1}, \alpha_j)$. Let ζ_j be the only zero of g on the interval (α_{j+1}, α_j) .

Proposition 5. *Let $a \neq b$. If $\delta \leq |\varphi_1(a)|$, then the extremal spectral point $(\widehat{\lambda}_1, \widehat{\lambda}_2)$ is unique, belongs to Λ_2 and is determined by the condition $\zeta_0 < \widehat{\lambda}_2 < \alpha_0$.*

Proposition 6. *Assume that $|\varphi_1(a)| < \delta < |\varphi_0(a)|$ and*

$$\gamma = \left| \frac{b-a}{1-ab} \right| \geq b^{2/3},$$

then the conclusion of Proposition 5 is valid, that is, the extremal spectral point $(\widehat{\lambda}_1, \widehat{\lambda}_2)$ is unique, belongs to Λ_2 and is determined by the condition $\zeta_0 < \widehat{\lambda}_2 < \alpha_0$.

5. OPTIMAL RECOVERY METHOD

To construct optimal recovery methods we need the following result (several results of this type may be found in [2], [1], [3]).

Theorem 5. *Assume that there exist $\widehat{\lambda}_j \geq 0, j = 1, \dots, n$, such that the value of the extremal problem*

$$\|f_0\|_{L^2(\mu_0)}^2 \rightarrow \max, \quad \sum_{j=1}^{\infty} \widehat{\lambda}_j \|f_j\|_{L^2(0, \mu_j)}^2 \leq \sum_{j=1}^{\infty} \widehat{\lambda}_j \delta_j^2, \quad f \in X,$$

is the same as in (1). Moreover, assume that for every $\tilde{y} = (\tilde{y}_1, \dots, \tilde{y}_n) \in Y_1 \times \dots \times Y_n$, where Y_j are dense in $L^2(\mu_j)$, there exists $f_{\tilde{y}}$ which is a solution of the extremal problem

$$\sum_{j=1}^{\infty} \widehat{\lambda}_j \|f_j - \tilde{y}_j\|_{L^2(0, \mu_j)}^2 \rightarrow \min, \quad f \in X.$$

Moreover, let $\widehat{A}: L^2(\mu_1) \times \dots \times L^2(\mu_n) \rightarrow L^2(\mu_0)$ be a linear continuous operator, where the norm in $L^2(\mu_1) \times \dots \times L^2(\mu_n)$ is defined as

$$\|y\| = \left(\sum_{j=1}^n \|y_j\|_{L^2(\mu_j)}^2 \right)^{1/2},$$

such that for all $\tilde{y} = (\tilde{y}_1, \dots, \tilde{y}_n) \in Y_1 \times \dots \times Y_n$,

$$\widehat{A}(\tilde{y}) = (f_{\tilde{y}})_0.$$

Then

$$E(X, \mathcal{D}, \mu, \delta) = \sup_{\substack{f \in X \\ \|f_j\|_{L^2(\mu_j)} \leq \delta_j, j=1, \dots, n}} \|f_0\|_{L^2(\mu_0)}$$

and the method $\widehat{A}(y)$ is optimal.

We will apply Theorem 5 to the construction of optimal recovery method for the Schwarz Lemma type problem considered above.

Consider the extremal problem

$$(4) \quad \int_{\Gamma} |f|^2 d\mu \rightarrow \max, \quad \widehat{\lambda}_1 |f(a)|^2 + \widehat{\lambda}_2 \|f\|_{H^2}^2 \leq \widehat{\lambda}_1 \delta^2 + \widehat{\lambda}_2, \quad f \in H^2,$$

where as before μ is the normalized Lebesgue measure on Γ and $(\widehat{\lambda}_1, \widehat{\lambda}_2)$ is an extremal spectral point for problem (2).

Proposition 7. *Suppose that either*

1. $a \neq b$ and $\delta \leq |\varphi_1(a)|$, or $|\varphi_1(a)| < \delta < |\varphi_0(a)|$ and $\gamma = \left| \frac{b-a}{1-ab} \right| \geq b^{2/3}$,

or

2. $a = b$ and $\delta < \varphi(b) = 1/\sqrt{1-b^2}$.

Then the values of extremal problems (2) and (4) are the same.

Theorem 6. *Suppose that one of the following conditions is satisfied*

1. $\delta \geq |\varphi_0(a)|$,
2. $\delta \leq |\varphi_1(a)|$,
3. $|\varphi_1(a)| < \delta < |\varphi_0(a)|$, $\gamma \geq b^{2/3}$,
4. $a = b$,

and $(\widehat{\lambda}_1, \widehat{\lambda}_2)$ is the corresponding extremal spectral point. Then the error of optimal recovery is given by

$$\sqrt{\widehat{\lambda}_1 \delta^2 + \widehat{\lambda}_2}$$

and the method

$$(5) \quad \widehat{A}(y)(z) = \frac{\widehat{\lambda}_1 y}{\widehat{\lambda}_1 + \widehat{\lambda}_2(1-|a|^2)} \cdot \frac{1-|a|^2}{1-\bar{a}z}$$

is optimal.

Note that for $a = b$ the optimal method of recovery (5) does not depend on δ and has the form

$$\widehat{A}(y)(z) = \frac{1-|b|^2}{1-bz}.$$

6. OPEN PROBLEMS

1. It would be desirable to identify the extremal spectral point in all possible cases. We have shown that in a number of cases the extremal spectral point is the only point in Λ_2 such that $\zeta_0 < \widehat{\lambda}_2 < \alpha_0$. Our attempts to find a nontrivial-case when this point is not extremal failed.

Thus, we are tempted to conjecture that the point of Λ_2 with the biggest λ_2 is always extremal.

Conjecture. *If $a \neq b$ and $\delta < |\varphi_0(a)|$, the point in Λ_2 such that $\zeta_0 < \widehat{\lambda}_2 < \alpha_0$ is always the spectral extremal point for problem (2).*

2. It is natural to ask which choice of a minimizes the value of problem (2) (of course, this choice of a leads to the least optimal recovery error). It follows from above discussion that the point b plays a special role.

Problem. *Does the choice $a = b$ always lead to the least mean square optimal recovery error?*

3. Finally, if in problem (2) we replace the constraint $|\varphi(a)| \leq \delta$ with

$$\frac{1}{2\pi r} \int_{|z-a|=r} |f(z)|^2 |d(z-a)| \leq \delta, \quad 0 < r < 1 - |a|,$$

then the problem becomes even more difficult. The reason is that in the right hand side of Euler's equation the term $\lambda_1 \frac{f(a)}{1 - \bar{a}z}$ is replaced with

$$\lambda_1 f \left(a - \frac{r^2 z}{1 - \bar{a}z} \right)$$

and the equation turns into

$$\frac{1}{1 - \rho w} f \left(\frac{\rho}{1 - \rho w} \right) = \frac{\lambda_1}{1 - \bar{a}z} f \left(a - \frac{r^2 z}{1 - \bar{a}z} \right) + \lambda_2 f(w).$$

Thus, finding the spectrum in this case is reduced to finding eigenvalues of an operator which is a linear combination of two compact non-commuting operators. It would be very interesting to find the eigenbasis which corresponds to this problem and to find the solution.

REFERENCES

- [1] G. G. Magaril-Il'yaev and K. Yu. Osipenko, "Optimal recovery of functions and their derivatives from inaccurate information about a spectrum and inequalities for derivatives", *Funkc. analiz i ego prilozh.* **37** (2003), 51–64; English transl. in *Funct. Anal and Its Appl.*, **37** (2003).
- [2] A. A. Melkman and C. A. Micchelli, "Optimal estimation of linear operators in Hilbert spaces from inaccurate data," *SIAM J. Numer. Anal.*, **16** (1979), 87–105.
- [3] K. Yu. Osipenko, "The Hardy–Littlewood–Pólya inequality for analytic functions from Hardy–Sobolev spaces", *Mat. Sb.* **197** (2006), 15–34; English transl. in *Sbornic: Mathematics* **197** (2006), 315–334.
- [4] K. Yu. Osipenko, M. I. Stessin, "On problems of recovery in Hardy and Bergman spaces", *Mat. Zametki* **49** (1991), 95–104; English transl. in *Math. Notes* **49** (1991), 395–401.

- [5] K. Yu. Osipenko, M. I. Stessin, "On optimal recovery of a holomorphic function in the unit ball of \mathbb{C}^n ", *Constr. Approx.* **8** (1992), 141–159.
- [6] K. Yu. Osipenko, M. I. Stessin, "On some problems of optimal recovery of analytic and harmonic functions from inaccurate data", *J. Approx. Theory* **70** (1992), 206–228.