

# OPTIMAL RECOVERY OF PERIODIC FUNCTIONS FROM FOURIER COEFFICIENTS GIVEN WITH AN ERROR

K. YU. OSIPENKO

ABSTRACT. We construct optimal methods of recovery of  $2\pi$ -periodic functions analytic in a strip and its derivatives at a point  $t \in [0, 2\pi)$ , using information about the Fourier coefficients given with an error in the uniform norm. The same problem is solved for the Sobolev space  $\widetilde{W}_2^r$ .

## 1. INTRODUCTION

Let  $X$  and  $Y$  be linear spaces over the field  $K = \mathbb{R}$  or  $\mathbb{C}$ ,  $W \subset X$  and  $U \subset Y$  balanced convex sets and  $I: W \rightarrow Y$  a linear operator. Denote by  $X'$  the set of all linear functionals on  $X$ . We consider the problem of optimal recovery of  $\langle x', x \rangle$  where  $x' \in X'$  and  $x \in W$ , using information about approximate values of the operator  $I$ . A method of recovery is any function  $\varphi: Y \rightarrow K$ . The value

$$(1) \quad e(x', I, W, U) := \inf_{\varphi} \sup_{x \in W} \sup_{\substack{y \in Y \\ Ix - y \in U}} |\langle x', x \rangle - \varphi(y)|$$

is called the intrinsic error in the recovery problem. Any  $\varphi_0$  for which

$$e(x', I, W, U) = \sup_{x \in W} \sup_{\substack{y \in Y \\ Ix - y \in U}} |\langle x', x \rangle - \varphi_0(y)|$$

is said to be an optimal method.

Many examples and other settings of optimal recovery problems can be found in [1]–[6]. It follows from Magaril-Il'yaev and Osipenko [6] that there is a linear optimal method  $\varphi_0(y) = \langle y', y \rangle$ ,  $y' \in Y'$ , and the following equality

$$(2) \quad e(x', I, W, U) = \sup_{\substack{x \in W \\ Ix \in U}} |\langle x', x \rangle|$$

---

The research was supported in part by Russian Foundation of Fundamental Research (Grant 93-01-00237) and by the International Scientific Fund (Grant MR 1000).

holds. On the other hand, since  $U$  and  $W$  are balanced, we have

$$(3) \quad e(x', I, W, U) = \inf_{y' \in Y'} \sup_{\substack{x \in W \\ z \in U}} | \langle x', x \rangle - \langle y', Ix - z \rangle | \\ = \inf_{y' \in Y'} \left( \sup_{x \in W} | \langle x', x \rangle - \langle y', Ix \rangle | + \sup_{z \in U} | \langle y', z \rangle | \right).$$

In this paper, we consider the problem of optimal recovery of  $2\pi$ -periodic functions analytic in a strip and its derivatives from the Hardy–Sobolev and Bergman–Sobolev spaces based on the information about Fourier coefficients given with an error in the uniform norm. We also obtain an optimal method of recovery in the analogous problem for the Sobolev space  $\widetilde{W}_2^r$ .

A similar problem for the estimation of functions in the  $L_2$ -norm was considered in Melkman and Micchelli [4]. The case when the  $l_2$ -norm is used to measure the error in the Fourier coefficients was analyzed by Micchelli and Rivlin [1]. In Boyanov [7] the problem of optimal recovery of periodic functions from the Sobolev space  $\widetilde{W}_q^r$ ,  $1 \leq q \leq \infty$ , was solved for the case when the Fourier coefficients are known exactly.

## 2. OPTIMAL RECOVERY IN HILBERT SPACES FROM INACCURATE FOURIER COEFFICIENTS

Let  $X$  be a Hilbert space and  $e_1, e_2, \dots$  a complete orthonormal system in  $X$ . For  $x \in X$  denote by  $x_j := (x, e_j)$  the Fourier coefficients of  $x$ . Consider the problem (1) for  $W = BX := \{x \in X : \|x\| \leq 1\}$ ,  $\langle x', x \rangle = (x, f)$ ,  $f \in X$ ,  $|f_j| > 0$ ,  $j = 1, 2, \dots$ ,  $Ix = (x_1, \dots, x_n)$  and

$$U = \{y = (y_1, \dots, y_n) : |y_j| \leq \delta_j, j = 1, \dots, n\}.$$

Thus we consider the problem of optimal recovery of the linear functional  $(x, f)$  from approximate Fourier coefficients  $(\tilde{x}_1, \dots, \tilde{x}_n)$  such that

$$|x_j - \tilde{x}_j| \leq \delta_j, \quad j = 1, \dots, n.$$

In this case the intrinsic error will be denoted by  $e(f, I, BX, \delta)$ .

For  $a \in \mathbb{R}$  put

$$a_+ := \begin{cases} a, & a > 0, \\ 0, & a \leq 0. \end{cases}$$

**Theorem 1.** *Let  $\lambda \in (0, \|f\|)$  be a solution of the equation*

$$(4) \quad \|f\|^2 - \sum_{j=1}^n (|f_j|^2 - \lambda^2 \delta_j^2)_+ - \lambda^2 = 0.$$

*Then*

$$(5) \quad (x, f) \approx \sum_{j=1}^n (1 - \lambda \delta_j |f_j|^{-1})_+ \overline{f_j} \tilde{x}_j$$

is an optimal method of recovery and

$$e(f, I, BX, \delta) = \lambda + \sum_{j=1}^n \delta_j (|f_j| - \lambda \delta_j)_+.$$

*Proof.* First we show that the equation (4) has a solution  $\lambda \in (0, \|f\|]$ . Denote by  $\varphi(\lambda)$  the function on the left hand side of (4). This function is continuous for all  $\lambda \geq 0$ . Moreover,

$$\varphi(0) = \|f\|^2 - \sum_{j=1}^n |f_j|^2 > 0.$$

Since  $\varphi(\|f\|) < 0$ , there exists a  $\lambda \in (0, \|f\|]$  which is a solution of (4).

For such  $\lambda$  consider the method (5). In view of (3) we have

$$\begin{aligned} e(f, I, BX, \delta) &\leq \sup_{x \in BX} \left| (x, f) - \sum_{j=1}^n (1 - \lambda \delta_j |f_j|^{-1})_+ \bar{f}_j x_j \right| \\ &+ \sum_{j=1}^n \delta_j |f_j| (1 - \lambda \delta_j |f_j|^{-1})_+ = \sup_{x \in BX} (x, f_\lambda) + \sum_{j=1}^n \delta_j (|f_j| - \lambda \delta_j)_+ \end{aligned}$$

where

$$(f_\lambda)_j = \begin{cases} f_j, & j \geq n+1, \\ f_j - f_j (1 - \lambda \delta_j |f_j|^{-1})_+, & 1 \leq j \leq n. \end{cases}$$

It can be easily shown that

$$\|f_\lambda\|^2 = \|f\|^2 - \sum_{j=1}^n (|f_j|^2 - \lambda^2 \delta_j^2)_+ = \lambda^2.$$

Consequently

$$e(f, I, BX, \delta) \leq \lambda + \sum_{j=1}^n \delta_j (|f_j| - \lambda \delta_j)_+.$$

Put

$$x_0 := \frac{f_\lambda}{\|f_\lambda\|} = \lambda^{-1} f_\lambda.$$

Let  $1 \leq j \leq n$ . If  $1 - \lambda \delta_j |f_j|^{-1} > 0$  then

$$|(x_0)_j| = \lambda^{-1} |(f_\lambda)_j| = \delta_j.$$

If  $1 - \lambda \delta_j |f_j|^{-1} \leq 0$  then

$$|(x_0)_j| = \lambda^{-1} |f_j| \leq \delta_j.$$

Thus  $Ix_0 \in U$ . Using (2) we obtain

$$\begin{aligned} e(f, I, BX, \delta) &\geq |(x_0, f)| = \lambda^{-1} \left( \|f\|^2 - \sum_{j=1}^n |f_j|^2 (1 - \lambda\delta_j |f_j|^{-1})_+ \right) \\ &= \lambda^{-1} \left( \|f\|^2 - \sum_{j=1}^n (|f_j| + \lambda\delta_j) (|f_j| - \lambda\delta_j)_+ + \lambda \sum_{j=1}^n \delta_j (|f_j| - \lambda\delta_j)_+ \right) \\ &= \lambda + \sum_{j=1}^n \delta_j (|f_j| - \lambda\delta_j)_+. \end{aligned}$$

This completes the proof of the theorem.  $\square$

Now let  $\delta_j = \delta\lambda_j$ ,  $\lambda_j > 0$ ,  $j = 1, \dots, n$ , and  $\delta \geq 0$ .

**Theorem 2.** *Suppose that*

$$|f_1|\lambda_1^{-1} \geq \dots \geq |f_n|\lambda_n^{-1}.$$

Set

$$\mu_k := \left( \sum_{j=1}^k \lambda_j^2 + |f_k|^{-2} \lambda_k^2 \sum_{j=k+1}^{\infty} |f_j|^2 \right)^{-1/2}, \quad k = 1, \dots, n,$$

$\mu_0 := +\infty$ ,  $\mu_{n+1} := 0$ , and  $\Delta_k := [\mu_{k+1}, \mu_k)$ ,  $k = 0, \dots, n$ . Then for  $\delta \in \Delta_k$ ,  $0 \leq k \leq n$ , the method

$$(x, f) \approx \sum_{j=1}^k \left( 1 - \delta \frac{\lambda_j}{|f_j|} \sqrt{\frac{\sum_{j=k+1}^{\infty} |f_j|^2}{1 - \delta^2 \sum_{j=1}^k \lambda_j^2}} \right) \bar{f}_j \tilde{x}_j$$

is optimal and

$$e(f, I, BX, \delta) = \sqrt{\sum_{j=k+1}^{\infty} |f_j|^2} \sqrt{1 - \delta^2 \sum_{j=1}^k \lambda_j^2 + \delta \sum_{j=1}^k \lambda_j |f_j|}.$$

*Proof.* The equation (4) now takes the following form

$$(6) \quad \|f\|^2 - \sum_{j=1}^n (|f_j|^2 - \lambda^2 \delta^2 \lambda_j^2)_+ - \lambda^2 = 0.$$

If  $\delta = 0$ , then the solution of (6) is evident and the theorem follows from Theorem 1 immediately. If  $\delta > 0$ , then (6) is equivalent to the equation

$$(7) \quad \frac{c^2}{\|f\|^2 - \sum_{j=1}^n (|f_j|^2 - c^2 \lambda_j^2)_+} = \delta^2$$

where  $c = \lambda\delta$ . Denote by  $\varphi(c)$  the function on the left hand side of (7). It is easy to show that  $\varphi(c)$  is monotonically increasing for  $c \geq 0$ . Furthermore,

$$\varphi(|f_k|\lambda_k^{-1}) = \mu_k^2, \quad k = 1, \dots, n.$$

Hence for  $\delta \in \Delta_k$ ,  $0 \leq k \leq n$ ,

$$c = \delta \sqrt{\frac{\sum_{j=k+1}^{\infty} |f_j|^2}{1 - \delta^2 \sum_{j=1}^k \lambda_j^2}}$$

is the solution of (7). Now the theorem follows from Theorem 1.  $\square$

For  $\lambda_1 = \dots = \lambda_n = 1$ , Theorem 2 was proved in [8] using more complicated arguments.

Denote by  $L$  the linear space of vectors  $x = (x_1, x_2, \dots)$ ,  $x_j \in \mathbb{C}$ , which satisfy the condition

$$\sum_{j=1}^{\infty} \gamma_j |x_j|^2 < \infty$$

where  $\gamma_1 \geq 0$  and  $\gamma_j > 0$ ,  $j > 1$ . Let  $x'$  be the linear functional on  $L$  defined by the following equality

$$\langle x', x \rangle := \sum_{j=1}^{\infty} x_j \bar{f}_j$$

where  $|f_j| > 0$ ,  $j > 1$ , and

$$\sum_{j=2}^{\infty} \gamma_j^{-1} |f_j|^2 < \infty.$$

Consider the problem of optimal recovery of the functional  $x'$  on the set  $BL := \{x \in L : \sum_{j=1}^{\infty} \gamma_j |x_j|^2 \leq 1\}$  from information  $(\tilde{x}_1, \dots, \tilde{x}_n)$  such that

$$|x_j - \tilde{x}_j| \leq \delta \lambda_j, \quad \lambda_j > 0, \quad j = 1, \dots, n.$$

Put

$$m := \begin{cases} 1, & \gamma_1 f_1 \neq 0, \\ 2, & \gamma_1 f_1 = 0. \end{cases}$$

**Theorem 3.** *Suppose that*

$$(8) \quad \frac{|f_m|}{\lambda_m \gamma_m} \geq \dots \geq \frac{|f_n|}{\lambda_n \gamma_n}.$$

Set

$$\mu_{km} := \left( \sum_{j=m}^k \gamma_j \lambda_j^2 + \gamma_k^2 |f_k|^{-2} \lambda_k^2 \sum_{j=k+1}^{\infty} \gamma_j^{-1} |f_j|^2 \right)^{-1/2}, \quad k = m, \dots, n,$$

$\mu_{m-1, m} := +\infty$ ,  $\mu_{n+1, m} := 0$ , and  $\Delta_{km} := [\mu_{k+1, m}, \mu_{km}]$ ,  $k = m - 1, \dots, n$ . Then for  $\delta \in \Delta_{km}$ ,  $m - 1 \leq k \leq n$ , the method

$$(9) \quad \langle x', x \rangle \approx (m-1) \bar{f}_1 \tilde{x}_1 + \sum_{j=m}^k \nu_{jm} \bar{f}_j \tilde{x}_j,$$

where

$$\nu_{jm} = 1 - \delta \frac{\gamma_j \lambda_j}{|f_j|} \sqrt{\frac{\sum_{j=k+1}^{\infty} \gamma_j^{-1} |f_j|^2}{1 - \delta^2 \sum_{j=m}^k \gamma_j \lambda_j^2}},$$

is optimal and

$$e(x', I, BL, \delta) = \sqrt{\sum_{j=k+1}^{\infty} \gamma_j^{-1} |f_j|^2} \sqrt{1 - \delta^2 \sum_{j=m}^k \gamma_j \lambda_j^2} + \delta \sum_{j=m}^k \lambda_j |f_j|.$$

*Proof.* Consider the case  $m = 1$ . Then  $L$  is a Hilbert space with the inner product

$$(x, y)_L := \sum_{j=1}^{\infty} \gamma_j x_j \bar{y}_j.$$

The vectors  $e_1, e_2, \dots$ ,

$$(e_j)_s := \begin{cases} 0, & s \neq j, \\ \gamma_j^{-1/2}, & s = j, \end{cases}$$

form a complete orthonormal basis in  $L$ . The Fourier coefficients of  $x$  are equal to  $(x, e_j) = \sqrt{\gamma_j} x_j$ . Now we can use Theorem 2 in which we have to replace  $\lambda_j$  and  $f_j$  by  $\gamma_j^{1/2} \lambda_j$  and  $\gamma_j^{-1/2} f_j$ , respectively.

Suppose that  $\gamma_1 = 0$ . Denote by  $L_0$  the space of all vectors  $x \in L$  for which  $x_1 = 0$ . The space  $L_0$  is a Hilbert space with the inner product

$$(x, y)_{L_0} := \sum_{j=2}^{\infty} \gamma_j x_j \bar{y}_j.$$

From Theorem 2 it follows that the method

$$\langle x', x \rangle \approx \sum_{j=2}^k \nu_{jm} \bar{f}_j \tilde{x}_j$$

is optimal for the set  $BL_0$ , and

$$e(x', I, BL_0, \delta) = \sqrt{\sum_{j=k+1}^{\infty} \gamma_j^{-1} |f_j|^2} \sqrt{1 - \delta^2 \sum_{j=2}^k \gamma_j \lambda_j^2} + \delta \sum_{j=2}^k \lambda_j |f_j|.$$

From (2) we have

$$(10) \quad e(x', I, BL, \delta) = \delta \lambda_1 |f_1| + e(x', I, BL_0, \delta).$$

On the other hand, from (3) it follows that for the method (9)

$$\begin{aligned} e(x', I, BL, \delta) &\leq \sup_{x \in BL_0} \left| \langle x', x \rangle - \sum_{j=2}^k \nu_{jm} \bar{f}_j x_j \right| + \delta \lambda_1 |f_1| \\ &\quad + \sup_{|z_j| \leq \delta \lambda_j} \left| \sum_{j=2}^k \nu_{jm} \bar{f}_j z_j \right| = \delta \lambda_1 |f_1| + e(x', I, BL_0, \delta). \end{aligned}$$

In view of (10) the method (9) is optimal for the set  $BL$ .

Now assume that  $f_1 = 0$ . Since from (2)

$$e(x', I, BL, \delta) = e(x', I, BL_0, \delta),$$

it suffices to construct an optimal method for the set  $BL_0$ . It can be immediately obtained from Theorem 2. The theorem is proved.  $\square$

### 3. OPTIMAL RECOVERY IN HARDY-SOBOLEV AND BERGMAN-SOBOLEV SPACES

Let  $W$  be a shift invariant class of sufficiently smooth and  $2\pi$ -periodic functions. Consider the problem of optimal recovery of  $f^{(s)}(t)$ ,  $t \in [0, 2\pi)$ ,  $f \in W$ , using information about the Fourier coefficients

$$c_k = \frac{1}{2\pi} \int_0^{2\pi} f(t) e^{-ikt} dt, \quad |k| \leq n,$$

given with error at most  $\delta$  in the uniform norm, i.e., by  $\tilde{c}_k$  such that

$$|c_k - \tilde{c}_k| \leq \delta, \quad |k| \leq n.$$

Denote by  $e_{ns}(W, \delta)$  the intrinsic error for this problem (from (2) it follows that it does not depend on  $n$ ).

Let  $\tilde{H}_{2,\beta}$  be the space of all  $2\pi$ -periodic functions analytic in the strip  $S_\beta := \{z \in \mathbb{C} : |\operatorname{Im} z| < \beta\}$  which satisfy the condition

$$\|f\|_{\tilde{H}_{2,\beta}} := \sup_{0 \leq \eta < \beta} \left( \frac{1}{4\pi} \int_0^{2\pi} (|f(t+i\eta)|^2 + |f(t-i\eta)|^2) dt \right)^{1/2} < \infty.$$

The Hardy-Sobolev space  $\tilde{H}_{2,\beta}^r$  is the set of all  $2\pi$ -periodic functions analytic in the strip  $S_\beta$  for which  $f^{(r)} \in \tilde{H}_{2,\beta}$ . Set

$$B\tilde{H}_{2,\beta}^r := \left\{ f \in \tilde{H}_{2,\beta}^r : \|f^{(r)}\|_{\tilde{H}_{2,\beta}} \leq 1 \right\}, \quad r = 0, 1, \dots$$

Functions from  $\tilde{H}_{2,\beta}$  have finite boundary values almost everywhere and the space  $\tilde{H}_{2,\beta}$  can be considered as a Hilbert space with the inner product

$$(f, g)_{\tilde{H}_{2,\beta}} := \frac{1}{4\pi} \int_0^{2\pi} \left( f(t+i\beta) \overline{g(t+i\beta)} + f(t-i\beta) \overline{g(t-i\beta)} \right) dt.$$

It is easy to verify that the functions  $e_j(z) := e^{ijz}$ ,  $j = 0, \pm 1, \dots$  form a complete orthogonal basis in  $\tilde{H}_{2,\beta}$  and  $\|e_j\|_{\tilde{H}_{2,\beta}}^2 = \cosh 2j\beta$ . Thus  $f \in B\tilde{H}_{2,\beta}^r$  iff

$$f(z) = \sum_{j=-\infty}^{+\infty} c_j e^{ijz}$$

and

$$\sum_{j=-\infty}^{+\infty} |c_j|^2 j^{2r} \cosh 2j\beta \leq 1.$$

For  $p = \{p_j\}_{-\infty}^{+\infty}$ ,  $p_j > 0$ , we introduce the following notation

$$\mu_{kr}(p, s) := \left( \sum_{|j| \leq k} j^{2r} p_j + k^{4r-2s} p_k^2 \sum_{|j| > k} j^{2(s-r)} p_j^{-1} \right)^{-1/2}, \quad 1 \leq k \leq n,$$

$$\mu_{n+1,r}(p, s) := 0, \quad \mu_{00}(p, s) := \left( \sum_{|j| \geq 0} p_j^{-1} \right)^{-1/2},$$

$$\mu_{0r}(p, s) := +\infty, \quad r \geq 1,$$

$$\Delta_{kr}(p, s) := [\mu_{k+1,r}(p, s), \mu_{kr}(p, s)], \quad 0 \leq k \leq n, \quad r \geq 0,$$

$$\Delta_{-1,0}(p, s) := [\mu_{00}(p, s), +\infty].$$

Using Theorem 3 with  $\bar{f}_j = (ij)^s e^{ijt}$ ,  $\lambda_j = 1$  and  $\gamma_j = j^{2r} \cosh 2j\beta$ , we obtain the following result.

**Theorem 4.** *Let  $r$  and  $s$  be nonnegative integers such that  $0 \leq s \leq 2r$ . Put  $p_j = \cosh 2j\beta$ ,  $j = 0, \pm 1, \dots$ . For  $\delta \in \Delta_{kr}(p, s)$  the method*

$$(11) \quad f^{(s)}(t) \approx \sum_{|j| \leq k} \nu_{jk}(p, s, \delta) \tilde{c}_j (ij)^s e^{ijt},$$

where

$$\nu_{jk}(p, s, \delta) = 1 - \delta |j|^{2r-s} p_j \sqrt{\frac{\sum_{|j| > k} j^{2(s-r)} p_j^{-1}}{1 - \delta^2 \sum_{|j| \leq k} j^{2r} p_j}},$$

is optimal for the class  $B\tilde{H}_{2,\beta}^r$ , and

$$e_{ns}(B\tilde{H}_{2,\beta}^r, \delta) = E_{kr}(p, s, \delta) := \sqrt{\sum_{|j| > k} j^{2(s-r)} p_j^{-1}} \sqrt{1 - \delta^2 \sum_{|j| \leq k} j^{2r} p_j} + \delta \sum_{|j| \leq k} |j|^s.$$

Denote by  $\tilde{A}_{2,\beta}$  the space of all  $2\pi$ -periodic functions analytic in the strip  $S_\beta$  which satisfy the condition

$$\|f\|_{\tilde{A}_{2,\beta}} := \left( \frac{1}{4\pi\beta} \int_0^{2\pi} \int_{-\beta}^{\beta} |f(t+i\eta)|^2 dt d\eta \right)^{1/2} < \infty.$$

The Bergman–Sobolev space  $\tilde{A}_{2,\beta}^r$  is the set of all  $2\pi$ -periodic functions analytic in the strip  $S_\beta$  for which  $f^{(r)} \in \tilde{A}_{2,\beta}$ . Set

$$B\tilde{A}_{2,\beta}^r := \left\{ f \in \tilde{A}_{2,\beta}^r : \|f^{(r)}\|_{\tilde{A}_{2,\beta}} \leq 1 \right\}, \quad r = 0, 1, \dots .$$

Consider the problem of optimal recovery of  $f^{(s)}(t)$  for the class  $B\tilde{A}_{2,\beta}^r$ .  $\tilde{A}_{2,\beta}$  is a Hilbert space with the inner product

$$(f, g)_{\tilde{A}_{2,\beta}} := \frac{1}{4\pi\beta} \int_0^{2\pi} \int_{-\beta}^{\beta} f(t+i\eta) \overline{g(t+i\eta)} dt d\eta.$$

It can be easily shown that the functions  $e_j(z)$ ,  $j = 0, \pm 1, \dots$  form a complete orthogonal basis in  $\tilde{A}_{2,\beta}$  and

$$\|e_0\|_{\tilde{A}_{2,\beta}} = 1, \quad \|e_j\|_{\tilde{A}_{2,\beta}}^2 = \frac{\sinh 2j\beta}{2j\beta}, \quad j = \pm 1, \pm 2, \dots .$$

Therefore  $f \in B\tilde{A}_{2,\beta}^r$  iff

$$f(z) = \sum_{j=-\infty}^{+\infty} c_j e^{ijz}$$

and

$$\sum_{j=-\infty}^{+\infty} |c_j|^2 j^{2r} \|e_j\|_{\tilde{A}_{2,\beta}}^2 \leq 1.$$

Analogously to Theorem 4 we have

**Theorem 5.** *Let  $0 \leq s \leq 2r$ . Put*

$$p_0 = 1, \quad p_j = \frac{\sinh 2j\beta}{2j\beta}, \quad j = \pm 1, \pm 2, \dots .$$

*For  $\delta \in \Delta_{kr}(p, s)$  the method (11) is an optimal method for the class  $B\tilde{A}_{2,\beta}^r$  and*

$$e_{ns}(B\tilde{A}_{2,\beta}^r, \delta) = E_{kr}(p, s, \delta).$$

*Remark.* We need the condition  $0 \leq s \leq 2r$  to satisfy (8). For  $s > 2r$  optimal methods of recovery for the classes  $B\tilde{H}_{2,\beta}^r$  and  $B\tilde{A}_{2,\beta}^r$  can be constructed by Theorem 1.

Almost the same arguments as in Theorem 4 and Theorem 5 enable us to obtain an optimal method of recovery of  $f^{(s)}(t)$ ,  $0 \leq s \leq r-1$ , for

the Sobolev class  $\widetilde{B\bar{W}}_2^r$  which is the set of all real-valued  $2\pi$ -periodic functions such that  $f^{(r-1)}$  is absolutely continuous and

$$\frac{1}{2\pi} \int_0^{2\pi} |f^{(r)}(t)|^2 dt \leq 1.$$

**Theorem 6.** *Let  $0 \leq s \leq r - 1$ . Put  $p_j = 1$ ,  $j = 0, \pm 1, \dots$ . For  $\delta \in \Delta_{kr}(p, s)$  the method (11) is an optimal method for the class  $\widetilde{B\bar{W}}_2^r$  and*

$$e_{ns}(\widetilde{B\bar{W}}_2^r, \delta) = E_{kr}(p, s, \delta).$$

#### REFERENCES

- [1] C. A. Micchelli and T. J. Rivlin, *A survey of optimal recovery*, Optimal Estimation in Approximation Theory (C. A. Micchelli and T. J. Rivlin, eds), Plenum Press, New York, 1977, pp. 1–54.
- [2] C. A. Micchelli and T. J. Rivlin, *Lectures on optimal recovery*, Lect. Notes Math. **1129** (1985), 21–93.
- [3] J. F. Traub and H. Woźniakowski, *A General Theory of Optimal Algorithms*, Academic Press, New York, 1980.
- [4] A. A. Melkman and C. A. Micchelli, *Optimal estimation of linear operators in Hilbert spaces from inaccurate data*, SIAM J. Numer. Anal. **16** (1979), 87–105.
- [5] J. F. Traub, G. W. Wasilkowski and H. Woźniakowski, *Information-Based Complexity*, Academic Press, New York, 1988.
- [6] G. G. Magaril-Il'yaev and K. Yu. Osipenko, *Optimal recovery of functionals based on inaccurate data*, Mat. Zametki, **50** (1991), 85–93; English transl. in Math. Notes **50** (1991), 1274–1279.
- [7] B. D. Bojanov, *Best reconstruction of differentiable periodic functions from their Fourier coefficients*, Serdica **12** (1976), 300–304. (Russian)
- [8] K. Yu. Osipenko, *Best approximation methods and the order of informativeness of systems*, Mat. Sb **111** (1980), 532–556; English transl. in Math. USSR Sb. **39** (1981), 479–500.

DEPARTMENT OF MATHEMATICS, MOSCOW STATE UNIVERSITY OF AVIATION TECHNOLOGY, MOSCOW, RUSSIA, 103767