

Recovery of Derivatives for Functions Defined on the Semiaxis

K. Yu. Osipenko¹

*Moscow State University,
Institute for Information Transmission Problems, Russian Academy of Sciences, Moscow*

Abstract

The paper is concerned with the recovery problem of derivatives at the origin from noisy information about functions defined on the semiaxis for the Sobolev class. The problem of S. B. Stechkin about approximation of derivatives by bounded linear functionals is also studied.

Keywords: optimal recovery, inequalities for derivatives, approximation by bounded linear functionals

2010 MSC: 41A65, 41A44, 41A17, 49N30

1. Setting of Problems

Let $W_2^n(\mathbb{R}_+)$, $n \in \mathbb{N}$, be the Sobolev space of functions $x(\cdot) \in L_2(\mathbb{R}_+)$ such that the $(n-1)$ -st derivative is locally absolutely continuous and $x^{(n)}(\cdot) \in L_2(\mathbb{R}_+)$. Set

$$W_2^n(\mathbb{R}_+) = \{x(\cdot) \in W_2^n(\mathbb{R}_+) : \|x^{(n)}(\cdot)\|_{L_2(\mathbb{R}_+)} \leq 1\}.$$

We consider the problem of recovery of $x^{(k)}(0)$, $0 \leq k < n$, on the class $W_2^n(\mathbb{R}_+)$ by inaccurate information about $x(\cdot)$. We assume that for every function $x(\cdot) \in W_2^n(\mathbb{R}_+)$ we know $y(\cdot) \in L_2(\mathbb{R}_+)$ such that

$$\|x(\cdot) - y(\cdot)\|_{L_2(\mathbb{R}_+)} \leq \delta, \quad \delta > 0.$$

For a given $y(\cdot)$ we have to construct an approximate value of $x^{(k)}(0)$.

As recovery methods we consider all possible mappings $m: L_2(\mathbb{R}_+) \rightarrow \mathbb{R}$. The error of a method m is defined as follows

$$e_k(W_2^n(\mathbb{R}_+), \delta, m) = \sup_{\substack{x(\cdot) \in W_2^n(\mathbb{R}_+), y(\cdot) \in L_2(\mathbb{R}_+) \\ \|x(\cdot) - y(\cdot)\|_{L_2(\mathbb{R}_+)} \leq \delta}} |x^{(k)}(0) - m(y(\cdot))|.$$

Email address: kosipenko@yahoo.com (K. Yu. Osipenko)

¹The research was carried out with the financial support of the Russian Foundation for Basic Research (grant no. 17-01-00649)

The quantity

$$E_k(W_2^n(\mathbb{R}_+), \delta) = \inf_{m: L_2(\mathbb{R}_+) \rightarrow \mathbb{R}} e_k(W_2^n(\mathbb{R}_+), \delta, m)$$

is known as the optimal recovery error and a method for which this infimum is attained is called optimal.

Note that optimal recovery problems are closely connected with statistical estimation problems. Details may be found in [2].

We also study the problem of best approximation of $x^{(k)}(0)$ on the class $W_2^n(\mathbb{R}_+)$ by linear continuous functionals on $L_2(\mathbb{R}_+)$ with the norm not greater than some fixed positive number N (this problem is known as Stechkin's problem). It relies on finding the value

$$S_k(W_2^n(\mathbb{R}_+), N) = \inf_{\substack{y^* \in L_2(\mathbb{R}_+) \\ \|y^*\|_{L_2(\mathbb{R}_+)} \leq N}} \sup_{x(\cdot) \in W_2^n(\mathbb{R}_+)} |x^{(k)}(0) - \langle y^*, x(\cdot) \rangle|,$$

and also a functional delivering the lower bound which is called extremal.

Solutions of the formulated problems are closely connected with the problems of exact constants in Kolmogorov-type inequalities for derivatives. For the semiaxis the corresponding results which we essentially use here were obtained in [3] and [4]. For \mathbb{R} the analogous problems of recovery and approximation by bounded linear functionals were considered in [6]. The range of problems connected with Stechkin's problem was elucidated in the survey paper [1].

2. Main Results

It may be obtained from [4] that there exists a function $\widehat{x}(\cdot) \in \mathcal{W}_2^{2n}(\mathbb{R}_+)$ such that for all $x(\cdot) \in \mathcal{W}_2^n(\mathbb{R}_+)$ the following equality

$$x^{(k)}(0) = \int_{\mathbb{R}_+} x(t) \overline{\widehat{x}(t)} dt + \int_{\mathbb{R}_+} x^{(n)}(t) \overline{\widehat{x}^{(n)}(t)} dt \quad (1)$$

holds. We give the explicit form of $\widehat{x}(\cdot)$ and prove the corresponding result for completeness.

Put

$$\lambda_j = e^{(n+2j-1)\frac{i\pi}{2n}}, \quad j = 1, \dots, n, \quad A = \begin{pmatrix} \lambda_1^n & \dots & \lambda_n^n \\ \dots & \dots & \dots \\ \lambda_1^{2n-1} & \dots & \lambda_n^{2n-1} \end{pmatrix},$$

and $A_{s,j}$ is the cofactor of λ_j^{n+s-1} . Denote by $|A|$ the determinant of matrix A .

Lemma 1. *Set*

$$\widehat{x}(t) = \frac{(-1)^{n-k}}{|A|} \sum_{j=1}^n A_{n-k,j} e^{\lambda_j t}.$$

Then for all $x(\cdot) \in \mathcal{W}_2^n(\mathbb{R}_+)$ (1) holds, moreover,

$$\widehat{x}^{(k)}(0) = \widehat{A}_{n,k}^2, \quad (2)$$

where

$$\widehat{A}_{n,k} = \frac{1}{\sin^{1/2}((2k+1)\alpha)} \prod_{j=1}^k \cot j\alpha, \quad \alpha = \frac{\pi}{2n}.$$

Proof. Since $\lambda_j^{2n} = (-1)^{n-1}$ for all $j = 1, \dots, n$ we have

$$\widehat{x}(t) + (-1)^n \widehat{x}^{(2n)}(t) = 0.$$

In view of the fact that $\operatorname{Re} \lambda_j < 0$ for all $j = 1, \dots, n$, $\widehat{x}(\cdot) \in \mathcal{W}_2^{2n}(\mathbb{R}_+)$. Thus, for every $x(\cdot) \in \mathcal{W}_2^n(\mathbb{R}_+)$ we have

$$\int_{\mathbb{R}_+} x(t) \overline{\widehat{x}(t)} dt + (-1)^n \int_{\mathbb{R}_+} x(t) \overline{\widehat{x}^{(2n)}(t)} dt = 0.$$

Using integration by parts we obtain

$$\begin{aligned} \int_{\mathbb{R}_+} x(t) \overline{\widehat{x}(t)} dt + (-1)^n \sum_{p=1}^n (-1)^p x^{(p-1)}(0) \overline{\widehat{x}^{(2n-p)}(0)} \\ + \int_{\mathbb{R}_+} x^{(n)}(t) \overline{\widehat{x}^{(n)}(t)} dt = 0. \end{aligned} \quad (3)$$

For all $s, p = 1, \dots, n$ we have

$$\sum_{j=1}^n A_{s,j} \lambda_j^{n+p-1} = \delta_{p,s} |A|,$$

where $\delta_{p,s}$ is the Kronecker delta. Consequently,

$$\widehat{x}^{(2n-p)}(0) = \frac{(-1)^{n-k}}{|A|} \sum_{j=1}^n A_{n-k,j} \lambda_j^{2n-p} = (-1)^{n-k} \delta_{n-p+1, n-k}. \quad (4)$$

Substituting (4) in (3) we obtain (1).

Now let us calculate $\widehat{x}^{(k)}(0)$. We have

$$\widehat{x}^{(k)}(0) = \frac{(-1)^{n-k}}{|A|} \sum_{j=1}^n A_{n-k,j} \lambda_j^k = (-1)^{n-k} \frac{|A_k|}{|A|},$$

where

$$A_k = \begin{pmatrix} \lambda_1^n & \dots & \lambda_n^n \\ \dots & \dots & \dots \\ \lambda_1^{2n-k-2} & \dots & \lambda_n^{2n-k-2} \\ \lambda_1^k & \dots & \lambda_n^k \\ \lambda_1^{2n-k} & \dots & \lambda_n^{2n-k} \\ \dots & \dots & \dots \\ \lambda_1^{2n-1} & \dots & \lambda_n^{2n-1} \end{pmatrix}.$$

Since

$$\lambda_j^m = \sigma_m \mu_m^{j-1},$$

where

$$|\sigma_m| = 1, \quad \mu_m = e^{i \frac{\pi m}{n}},$$

matrices A and A_k may be rewritten in the forms

$$A = \begin{pmatrix} \sigma_n & \cdots & \sigma_n \mu_n^{n-1} \\ \cdots & \cdots & \cdots \\ \sigma_{2n-1} & \cdots & \sigma_{2n-1} \mu_{2n-1}^{n-1} \end{pmatrix},$$

$$A_k = \begin{pmatrix} \sigma_n & \cdots & \sigma_n \mu_n^{n-1} \\ \cdots & \cdots & \cdots \\ \sigma_{2n-k-2} & \cdots & \sigma_{2n-k-2} \mu_{2n-k-2}^{n-1} \\ \sigma_k & \cdots & \sigma_k \mu_k^{n-1} \\ \sigma_{2n-k} & \cdots & \sigma_{2n-k} \mu_{2n-k}^{n-1} \\ \cdots & \cdots & \cdots \\ \sigma_{2n-1} & \cdots & \sigma_{2n-1} \mu_{2n-1}^{n-1} \end{pmatrix}.$$

If we put $x(\cdot) = \widehat{x}(\cdot)$ in (1), then we obtain

$$\widehat{x}^{(k)}(0) = \|\widehat{x}(\cdot)\|_{L_2(\mathbb{R}_+)}^2 + \|\widehat{x}^{(n)}(\cdot)\|_{L_2(\mathbb{R}_+)}^2. \quad (5)$$

Thus, $\widehat{x}^{(k)}(0) > 0$. Using this fact and the formula for the Vandermonde determinant we have

$$\widehat{x}^{(k)}(0) = \prod_{\substack{j=1 \\ j \neq n-k}}^n \frac{|\mu_k - \mu_{n+j-1}|}{|\mu_{2n-k-1} - \mu_{n+j-1}|}.$$

Since

$$|\mu_p - \mu_s| = 2|\sin(p-s)\alpha|,$$

we obtain

$$\begin{aligned} \widehat{x}^{(k)}(0) &= \prod_{\substack{j=1 \\ j \neq n-k}}^n \frac{|\sin(n+j-k-1)\alpha|}{|\sin(n+j+k)\alpha|} \\ &= \frac{1}{\sin((2k+1)\alpha)} \frac{\prod_{j=k+1}^{k+n} \sin j\alpha}{\prod_{j=1}^k \sin j\alpha \prod_{j=1}^{n-k-1} \sin j\alpha} \\ &= \frac{1}{\sin((2k+1)\alpha)} \frac{\prod_{j=k+1}^{n-1} \sin j\alpha \prod_{j=n+1}^{n+k} \sin j\alpha}{\prod_{j=1}^k \sin j\alpha \prod_{j=1}^{n-k-1} \sin j\alpha} \\ &= \frac{1}{\sin((2k+1)\alpha)} \prod_{j=1}^k \cot j\alpha \frac{\prod_{j=k+1}^{n-1} \sin j\alpha}{\prod_{j=1}^{n-k-1} \sin j\alpha} \\ &= \frac{1}{\sin((2k+1)\alpha)} \prod_{j=1}^k \cot j\alpha \prod_{j=1}^{n-k-1} \cot j\alpha = \frac{1}{\sin((2k+1)\alpha)} \prod_{j=1}^k \cot^2 j\alpha. \end{aligned}$$

□

Theorem 1. *The following equality*

$$E_k(W_2^n(\mathbb{R}_+), \delta) = \widehat{A}_{n,k} \left(\frac{2n}{2n-2k-1} \right)^{\frac{2n-2k-1}{4n}} \left(\frac{2n}{2k+1} \right)^{\frac{2k+1}{4n}} \delta^{\frac{2n-2k-1}{2n}}$$

holds. Moreover,

$$\widehat{m}(y) = \beta^{k+1} \int_{\mathbb{R}_+} y(t) \overline{\widehat{x}(\beta t)} dt, \quad (6)$$

where

$$\beta = \left(\frac{2n-2k-1}{2k+1} \right)^{\frac{1}{2n}} \delta^{-\frac{1}{n}},$$

is the optimal method of recovery.

Proof. From (1) by the Cauchy-Schwarz inequality we obtain

$$\begin{aligned} |x^{(k)}(0)| &\leq \left(\|\widehat{x}(\cdot)\|_{L_2(\mathbb{R}_+)}^2 + \|\widehat{x}^{(n)}(\cdot)\|_{L_2(\mathbb{R}_+)}^2 \right)^{1/2} \\ &\quad \times \left(\|x(\cdot)\|_{L_2(\mathbb{R}_+)}^2 + \|x^{(n)}(\cdot)\|_{L_2(\mathbb{R}_+)}^2 \right)^{1/2}. \end{aligned}$$

Taking into account (5) we have

$$|x^{(k)}(0)| \leq \widehat{A}_{n,k} \left(\|x(\cdot)\|_{L_2(\mathbb{R}_+)}^2 + \|x^{(n)}(\cdot)\|_{L_2(\mathbb{R}_+)}^2 \right)^{1/2}. \quad (7)$$

Put $y(t) = \widehat{x}(bt)$, $b > 0$. Then

$$\|y(\cdot)\|_{L_2(\mathbb{R}_+)}^2 = \frac{1}{b} \|\widehat{x}(\cdot)\|_{L_2(\mathbb{R}_+)}^2, \quad \|y^{(n)}(\cdot)\|_{L_2(\mathbb{R}_+)}^2 = b^{2n-1} \|\widehat{x}^{(n)}(\cdot)\|_{L_2(\mathbb{R}_+)}^2.$$

Substituting $y(\cdot)$ into (7) we obtain that for all $b > 0$ the inequality

$$b^k \widehat{A}_{n,k}^2 \leq \widehat{A}_{n,k} \left(\frac{1}{b} \|\widehat{x}(\cdot)\|_{L_2(\mathbb{R}_+)}^2 + b^{2n-1} \|\widehat{x}^{(n)}(\cdot)\|_{L_2(\mathbb{R}_+)}^2 \right)^{1/2}$$

holds. In view of (5) and (2) we get that for all $b > 0$ the inequality $f(b) \geq 0$ is fulfilled, where

$$f(b) = b^{2n} \|\widehat{x}^{(n)}(\cdot)\|_{L_2(\mathbb{R}_+)}^2 - \widehat{A}_{n,k}^2 b^{2k+1} + \widehat{A}_{n,k}^2 - \|\widehat{x}(\cdot)\|_{L_2(\mathbb{R}_+)}^2.$$

It is easily seen that $f(\cdot)$ has the unique minimum on \mathbb{R}_+

$$b_0 = \left(\frac{(2k+1) \widehat{A}_{n,k}^2}{2n \|\widehat{x}^{(n)}(\cdot)\|_{L_2(\mathbb{R}_+)}^2} \right)^{\frac{1}{2n-2k-1}}.$$

On the other hand, $f(1) = 0$. Consequently, $b_0 = 1$. Thus,

$$\|\widehat{x}^{(n)}(\cdot)\|_{L_2(\mathbb{R}_+)} = \widehat{A}_{n,k} \sqrt{\frac{2k+1}{2n}}.$$

It follows from (5) and (2) that

$$\|\widehat{x}(\cdot)\|_{L_2(\mathbb{R}_+)} = \widehat{A}_{n,k} \sqrt{\frac{2n-2k-1}{2n}}.$$

Put $\widehat{x}_1(t) = \alpha \widehat{x}(\beta t)$, $\alpha, \beta > 0$. Choose α and β such that $\|\widehat{x}_1(\cdot)\|_{L_2(\mathbb{R}_+)} = \delta$ and $\|\widehat{x}_1^{(n)}(\cdot)\|_{L_2(\mathbb{R}_+)} = 1$. We have

$$\alpha^2 \beta^{-1} \|\widehat{x}(\cdot)\|_{L_2(\mathbb{R}_+)}^2 = \delta^2, \quad \alpha^2 \beta^{2n-1} \|\widehat{x}^{(n)}(\cdot)\|_{L_2(\mathbb{R}_+)}^2 = 1.$$

Hence,

$$\begin{aligned} \alpha &= \widehat{A}_{n,k}^{-1} \sqrt{\frac{2n}{2n-2k-1}} \left(\frac{2n-2k-1}{2k+1} \right)^{\frac{1}{4n}} \delta^{1-\frac{1}{2n}}, \\ \beta &= \left(\frac{2n-2k-1}{2k+1} \right)^{\frac{1}{2n}} \delta^{-\frac{1}{n}}. \end{aligned}$$

Substituting $\widehat{x}(t) = \alpha^{-1} \widehat{x}_1(t/\beta)$ in (1) we obtain

$$x^{(k)}(0) = \alpha^{-1} \int_{\mathbb{R}_+} x(t) \overline{\widehat{x}_1(t/\beta)} dt + \alpha^{-1} \beta^{-n} \int_{\mathbb{R}_+} x^{(n)}(t) \overline{\widehat{x}_1^{(n)}(t/\beta)} dt.$$

We change variables $t = \beta s$ and put $z(s) = x(\beta s)$. Then we have that for all $z(\cdot) \in \mathcal{W}_2^n(\mathbb{R}_+)$ the equality

$$z^{(k)}(0) = \lambda_1 \int_{\mathbb{R}_+} z(s) \overline{\widehat{x}_1(s)} ds + \lambda_2 \int_{\mathbb{R}_+} z^{(n)}(s) \overline{\widehat{x}_1^{(n)}(s)} ds \quad (8)$$

holds with

$$\lambda_1 = \frac{\beta^{k+1}}{\alpha}, \quad \lambda_2 = \frac{1}{\alpha \beta^{2n-k-1}}.$$

It follows from general results about optimal recovery of linear functionals (see, for example, [5]) that

$$E_k(W_2^n(\mathbb{R}_+), \delta) = \sup_{\substack{z(\cdot) \in W_2^n(\mathbb{R}_+) \\ \|z(\cdot)\|_{L_2(\mathbb{R}_+)} \leq \delta}} |z^{(k)}(0)|. \quad (9)$$

From (8) by the Cauchy-Schwarz inequality we obtain

$$E_k(W_2^n(\mathbb{R}_+), \delta) \leq \lambda_1 \delta^2 + \lambda_2. \quad (10)$$

Let us estimate the error of method (6), which may be written in the following way

$$\widehat{m}(y) = \lambda_1 \int_{\mathbb{R}_+} y(t) \overline{\widehat{x}_1(t)} dt.$$

Suppose that $z(\cdot) \in W_2^n(\mathbb{R}_+)$ and $\|z(\cdot) - y(\cdot)\|_{L_2(\mathbb{R}_+)} \leq \delta$. Taking into account (8) we have

$$\begin{aligned} & |z^{(k)}(0) - \widehat{m}(y)| \\ &= \left| z^{(k)}(0) - \lambda_1 \int_{\mathbb{R}_+} z(t) \overline{\widehat{x}_1(t)} dt + \lambda_1 \int_{\mathbb{R}_+} (z(t) - y(t)) \overline{\widehat{x}_1(t)} dt \right| \\ &= \left| \lambda_1 \int_{\mathbb{R}_+} (z(t) - y(t)) \overline{\widehat{x}_1(t)} dt + \lambda_2 \int_{\mathbb{R}_+} z^{(n)}(t) \overline{\widehat{x}_1^{(n)}(t)} dt \right| \leq \lambda_1 \delta^2 + \lambda_2. \end{aligned}$$

Consequently,

$$E_k(W_2^n(\mathbb{R}_+), \delta) \leq e_k(W_2^n(\mathbb{R}_+), \delta, \widehat{m}) \leq \lambda_1 \delta^2 + \lambda_2. \quad (11)$$

The last inequality together with (10) gives

$$\begin{aligned} E_k(W_2^n(\mathbb{R}_+), \delta) &= \lambda_1 \delta^2 + \lambda_2 \\ &= \widehat{A}_{n,k} \left(\frac{2n}{2n-2k-1} \right)^{\frac{2n-2k-1}{4n}} \left(\frac{2n}{2k+1} \right)^{\frac{2k+1}{4n}} \delta^{\frac{2n-2k-1}{2n}}. \end{aligned}$$

Inequality (11) implies also that \widehat{m} is the optimal method of recovery. \square

Note that the exact solution of extremal problem (9) gives us the exact inequality

$$|x^{(k)}(0)| \leq K_{nk} \|x(\cdot)\|_{L_2(\mathbb{R}_+)}^{\frac{2n-2k-1}{2n}} \|x^{(n)}(\cdot)\|_{L_2(\mathbb{R}_+)}^{\frac{2k+1}{2n}},$$

where

$$K_{nk} = \widehat{A}_{n,k} \left(\frac{2n}{2n-2k-1} \right)^{\frac{2n-2k-1}{4n}} \left(\frac{2n}{2k+1} \right)^{\frac{2k+1}{4n}}.$$

It may be also obtained from exact inequality (7) by Proposition 4 from [7, p. 119].

We now proceed to the Stechkin problem.

Theorem 2. *The following equality*

$$S_k(W_2^n(\mathbb{R}_+), N) = \widehat{A}_{n,k}^{\frac{2n}{2k+1}} \sqrt{\frac{2k+1}{2n-2k-1}} \left(\frac{2n-2k-1}{2n} \right)^{\frac{n}{2k+1}} N^{-\frac{2n-2k-1}{2k+1}}$$

holds. The functional

$$\langle \widehat{y}^*, x(\cdot) \rangle = \beta_N^{k+1} \int_{\mathbb{R}_+} x(t) \overline{\widehat{x}(\beta_N t)} dt$$

where

$$\beta_N = \left(\frac{2n}{2n-2k-1} \right)^{\frac{1}{2k+1}} \left(\frac{N}{\widehat{A}_{n,k}} \right)^{\frac{2}{2k+1}}$$

is extremal.

Proof. As was proved in the optimal recovery problem among all optimal methods there exists a method defined by a linear continuous functional, therefore

$$\begin{aligned} E_k(W_2^n(\mathbb{R}_+), \delta) &= \inf_{N>0} \inf_{\|y^*\|_{L_2(\mathbb{R}_+)} \leq N} \sup_{\substack{x(\cdot) \in W_2^n(\mathbb{R}_+), y(\cdot) \in L_2(\mathbb{R}_+) \\ \|x(\cdot) - y(\cdot)\|_{L_2(\mathbb{R}_+)} \leq \delta}} |x^{(k)}(0) - \langle y^*, y(\cdot) \rangle| \\ &\leq \inf_{\|y^*\|_{L_2(\mathbb{R}_+)} \leq N} \sup_{x(\cdot) \in W_2^n(\mathbb{R}_+)} |x^{(k)}(0) - \langle y^*, x(\cdot) \rangle| + \delta N = S_k(W_2^n(\mathbb{R}_+), N) + \delta N. \end{aligned}$$

Consequently, for all $N > 0$

$$S_k(W_2^n(\mathbb{R}_+), N) \geq E_k(W_2^n(\mathbb{R}_+), \delta) - \delta N. \quad (12)$$

We define the linear functional \widehat{y}^* as follows

$$\langle \widehat{y}^*, x(\cdot) \rangle = \lambda_1 \int_{\mathbb{R}_+} x(t) \widehat{x}_1(t) dt.$$

Then $\|\widehat{y}^*\|_{L_2(\mathbb{R}_+)} = \lambda_1 \delta$. If we choose δ such that $N = \lambda_1 \delta$, then it follows from (12) that

$$S_k(W_2^n(\mathbb{R}_+), N) \geq \lambda_2.$$

On the other hand, in view of (8) we have

$$S_k(W_2^n(\mathbb{R}_+), N) \leq \sup_{x(\cdot) \in W_2^n(\mathbb{R}_+)} |x^{(k)}(0) - \langle \widehat{y}^*, x(\cdot) \rangle| = \lambda_2.$$

Consequently, $S_k(W_2^n(\mathbb{R}_+), N) = \lambda_2$. If $N = \lambda_1 \delta$, then

$$\delta = \delta_N = \sqrt{\frac{2n-2k-1}{2k+1}} \left(\frac{2n-2k-1}{2n} \right)^{\frac{n}{2k+1}} \left(\frac{\widehat{A}_{n,k}}{N} \right)^{\frac{2n}{2k+1}}.$$

For $\delta = \delta_N$ we have

$$\lambda_2 = \widehat{A}_{n,k}^{\frac{2n}{2k+1}} \sqrt{\frac{2k+1}{2n-2k-1}} \left(\frac{2n-2k-1}{2n} \right)^{\frac{n}{2k+1}} N^{-\frac{2n-2k-1}{2k+1}}.$$

The functional \widehat{y}^* may be written in the following way

$$\langle \widehat{y}^*, x(\cdot) \rangle = \beta^{k+1} \int_{\mathbb{R}_+} x(t) \widehat{x}(\beta t) dt$$

where β is defined in Theorem 1. For $\delta = \delta_N$

$$\beta = \beta_N = \left(\frac{2n}{2n-2k-1} \right)^{\frac{1}{2k+1}} \left(\frac{N}{\widehat{A}_{n,k}} \right)^{\frac{2}{2k+1}}.$$

□

- [1] Arestov V. V. Approximation of unbounded operators by bounded operators and related extremal problems, *Russian Math. Surveys*, 51:6 (1996), 1093–1126.
- [2] Donoho D. L. Statistical estimation and optimal recovery, *The Annals of Statistics*, 22:1 (1994), 238–270.
- [3] Kalyabin G. A. Sharp constants in inequalities for intermediate derivatives (the Gabushin Case), *Funct. Anal. Appl.* 38:3 (2004), 184–191.
- [4] Lunev A. A., Oridoroga L. L. Exact constants in generalized inequalities for intermediate derivatives, *Math. Notes* 85:5 (2009), 703–711.
- [5] Magaril-Il'yaev G. G., Osipenko K. Yu. Optimal recovery of functionals based on inaccurate data, *Math. Notes* 50:6 (1991), 1274–1279.
- [6] Magaril-Il'yaev G. G., Osipenko K. Yu. Optimal recovery of values of functions and their derivatives from inaccurate data on the Fourier transform, *Sbornic: Mathematics* 195:10 (2004), 1461–1476.
- [7] Tikhomirov V. M. *Some Questions in Approximation Theory*, Moscow State University, Moscow, 1976 (in Russian).