

OPTIMAL RECOVERY OF FUNCTIONS IN $H_{2,\beta}$

K. YU. OSIPENKO (MOSCOW, RUSSIA)

Let $S_\beta := \{z \in \mathbb{C} : |\operatorname{Im} z| < \beta\}$ be a strip in the complex plane. Denote by $H_{2,\beta}$ the class of 2π -periodic, analytic in S_β functions f , which satisfy

$$\sup_{0 \leq \eta < \beta} \frac{1}{4\pi} \int_0^{2\pi} (|f(t + i\eta)|^2 + |f(t - i\eta)|^2) dt \leq 1.$$

Set

$$s_n(H_{2,\beta}, C(\mathbb{T})) := \inf_{t_1, \dots, t_n \in \mathbb{T}} \inf_{A: \mathbb{C}^n \rightarrow C(\mathbb{T})} \sup_{f \in H_{2,\beta}} \|f - A(f(t_1), \dots, f(t_n))\|_{C(\mathbb{T})},$$

$$i_n(H_{2,\beta}, C(\mathbb{T})) := \inf_{l_1, \dots, l_n} \inf_{A: \mathbb{C}^n \rightarrow C(\mathbb{T})} \sup_{f \in H_{2,\beta}} \|f - S(l_1 f, \dots, l_n f)\|_{C(\mathbb{T})},$$

where l_1, \dots, l_n are linear continuous functionals. The values $i_n(H_{2,\beta}, C(\mathbb{T}))$ are coincide with linear and Gel'fand n -widths which were calculated in [1]. We study the values $s_n(H_{2,\beta}, C(\mathbb{T}))$ and compare them with $i_n(H_{2,\beta}, C(\mathbb{T}))$.

Denote by K and K' be the complete elliptic integrals of the first kind with moduli k and $k' = \sqrt{1 - k^2}$. Suppose that k is defined by the equation

$$\frac{\pi K'}{2K} = \beta.$$

We prove that

$$\frac{s_{2n-1}(H_{2,\beta}, C(\mathbb{T}))}{i_{2n-1}(H_{2,\beta}, C(\mathbb{T}))} = 2\sqrt{\frac{kK}{\pi} \sinh \beta} + O(e^{-4\beta n}),$$

$$\frac{s_{2n}(H_{2,\beta}, C(\mathbb{T}))}{i_{2n}(H_{2,\beta}, C(\mathbb{T}))} = 2\sqrt{\frac{K}{\pi} \tanh \beta} + O(e^{-4\beta n}).$$

References

- [1] K. YU. OSIPENKO. On n -widths of holomorphic functions of several variables. J. Approx. Theory **82**, 1995, 135-155.
- [2] K. YU. OSIPENKO AND K. WILDEROTTER. Optimal information for approximating periodic analytic functions. Math. Comput. (to appear).